

MICRONUTRIENT DEFICIENCY INFORMATION SYSTEM

WORLD HEALTH ORGANIZATION

MDIS WORKING PAPER #2

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY



World Health Organization



United Nations Children's Fund

THE MICRONUTRIENT DEFICIENCY INFORMATION SYSTEM

The Micronutrient Deficiency Information System (MDIS) was established in 1991 in the Nutrition unit of the World Health Organization (WHO) in collaboration with the Community Systems Foundation of Ann Arbor, Michigan. The MDIS is a global surveillance mechanism for continually assessing the magnitude and distribution of deficiencies in three major micronutrients: iodine, vitamin A and iron. The databases provide the information required both to estimate the prevalence of these forms of micronutrient malnutrition on a national and global scale, and to provide timely and direct support for implementing and monitoring related prevention and control programmes.

Information is based on clinical indicators and selected biochemical and ecological parameters; it has been gathered from scientific journals, government documents, conference reports, and unpublished papers. To facilitate interpretation, every effort is made to specify the methods used in collecting and analysing information. Whenever possible, data are presented in tabular form stratified by administrative region, age, sex and nutrient status. On this basis, national "at-risk" and "affected" populations have been calculated. This degree of detail is provided in support of continuing efforts to standardize methodologies for assessing population-based deficiency rates. It is hoped that this will lead eventually both to improved monitoring of control activities and a better understanding of their impact within countries.

This is the second in the MDIS working paper series, available data on global prevalence of iodine deficiency disorders having been presented in 1993. A third report is in preparation, on the global prevalence of iron deficiency anaemia in children. Although previous efforts have been made to document the worldwide magnitude and distribution of micronutrient deficiencies, the MDIS is the most systematic and comprehensive approach developed thus far to providing robust epidemiological prevalence estimates of deficiencies of these three important micronutrients.

Because of the dynamic nature of global micronutrient malnutrition, WHO expects to update periodically this and other documents in the MDIS working paper series. Readers are urged to provide any missing information so that current estimates may be revised.


WHO/NUT/95.3
Distr.: General
Original: English

MICRONUTRIENT DEFICIENCY INFORMATION SYSTEM

WORLD HEALTH ORGANIZATION

MDIS WORKING PAPER #2

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY



World Health Organization



United Nations Children's Fund

©World Health Organization 1995

This document is not a formal publication of the World Health Organization (WHO), and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, reproduced or translated, in part or in whole, but not for sale or for use in conjunction with commercial purposes.

The views expressed in documents by named authors are solely the responsibility of those authors.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization welcomes requests for permission to reproduce or translate its documents, in part or in full. Applications and enquiries should be addressed to the Nutrition unit.

Micronutrient Deficiency Information System

World Health Organization
Nutrition unit
1211 Geneva 27 SWITZERLAND
Tel: 41.22.791.3318
Fax: 41.22.791.4156

Cover photo courtesy of the International Center for Eye Health

ACKNOWLEDGEMENTS

This document was prepared by the Nutrition unit, World Health Organization, Geneva with the technical support of the Micronutrient Deficiency Information System project initially based in Ann Arbor, Michigan, USA. Financial support was provided by UNICEF, New York and the Micronutrient Initiative, Ottawa, Canada. Dr Graeme Clugston heads the Nutrition unit and Dr Barbara Underwood is responsible there for matters dealing with micronutrient malnutrition. The group in Michigan is under the direction of Jonathan Gorstein and includes Suzinne Pak, Sanjaya, Peter Rycus, Helen Matzger and Claudia Gras. Special thanks are due to Suki McClatchey, seconded by James Akre of the Nutrition unit, for the excellent technical support she provided in organizing and presenting the information. Technical comments and review of sections of the document that deal with at-risk and affected populations were made by the following:

Dr David Alnwick	Senior Micronutrient Advisor, UNICEF
Dr Ken Bailey	Consultant, Nutrition unit, WHO
Dr Djamil Benbouzid	Nutrition unit, WHO
Dr Indira Chakravarty	Former Acting Regional Advisor, Nutrition unit, WHO Regional Office for South-East Asia, New Dehli
Dr Ian Darnton-Hill	Former Regional Advisor, Nutrition unit, WHO Regional Office for the Western Pacific, Manila
Dr Bruno de Benoist	Regional Officer, Nutrition unit, WHO Regional Office for Africa, Brazzaville
Dr Miguel Gueri	Regional Advisor, Nutrition unit, WHO Regional Office for the Americas, Washington, D.C.
Dr Elisabet Helsing	Regional Advisor, Nutrition unit, WHO Regional Office for Europe, Copenhagen
Dr John Mason	Technical Secretary, ACC/SCN
Dr Anna Verster	Regional Advisor, Nutrition unit, WHO Regional Office for the Eastern Mediterranean, Alexandria

TABLE OF CONTENTS

Preface	ix
-------------------	----

INTRODUCTION & BACKGROUND

Vitamin A deficiency (VAD)	1
Epidemiology of VAD	4
Ecological factors	5
Social factors	5
Economic factors	5
Clustering	6
Host factors	6
Age	6
Sex	6
Feeding practices	7
Disease patterns	7
Periods of increased physiological need	7
Indicators of VAD	8
Clinical indicators	8
Serum retinol	8
Other biological indicators	10

TABLES SUMMARIZING VAD PREVALENCE BY COUNTRY IN EACH WHO REGION

Interpreting data in the MDIS	15
Survey design	15
Subject selection	15
Data aggregation	15
Limitations	15
Guide to using summary tables	16
Summary of VAD status	
Global distribution of VAD prevalence	19
Regional distribution of VAD prevalence	
Africa	21

Americas	31
South-East Asia	39
Europe	45
Eastern Mediterranean	51
Western Pacific	57

DISAGGREGATED DATA TABLES

Guidelines for using disaggregated tables	65
Prevalence of ocular signs and symptoms	67
Prevalence of serum retinol levels suggestive of a public health problem	85

REFERENCES

References	97
------------------	----

ANNEXES

WHO regions	115
UNICEF regions	116

LIST OF TABLES

1. Estimated requirements for vitamin A ($\mu\text{g RE/day}$)	2
2. Available supply of vitamin A by WHO and UNICEF region ($\mu\text{g RE/day}$)	3
3. Type of data available by WHO region	4
4. Estimates of clinically affected and at-risk populations	4
5. Classification of xerophthalmia and prevalence criteria constituting a public health problem	9
6. Prevalence of VAD in children ≥ 1 year of age of serum values $\leq 0.70 \mu\text{mol/l}$	10

LIST OF ABBREVIATIONS

ACC/SCN	United Nations Administrative Committee on Coordination/Subcommittee on Nutrition
AED	Academy for Educational Development
AMR	Region of the Americas
AFR	African Region
CIC	Conjunctival Impression Cytology
EMR	Eastern Mediterranean Region
EUR	European Region
FAO	Food and Agriculture Organization of the United Nations
HKI	Helen Keller International
HPLC	High Pressure Liquid Chromatography
ICN	International Conference on Nutrition
IDA	Iron deficiency anaemia
IDD	Iodine deficiency disorders
IEC	Information, education and communication
IVACG	International Vitamin A Consultative Group
MDIS	Micronutrient Deficiency Information System
MOH	Ministry of health
MSG	Monosodium glutamate
PEM	Protein-energy malnutrition
RDA	Recommended dietary allowance
RDR	Relative dose response
RE	Retinol equivalents
SC	Save the Children
SEAR	South-East Asia Region
TFNC	Tanzania Food and Nutrition Centre
UNICEF	United Nations Children's Fund
USAID	United States Agency for International Development
VAD	Vitamin A deficiency
VAST	Vitamin A Supplementation Trials
VITAL	Vitamin A Field Support Project
WHO	World Health Organization
WHO/PAHO	World Health Organization/Pan American Health Organization
WPR	Western Pacific Region

PREFACE

Vitamin A deficiency (VAD), is the single most important cause of childhood blindness in developing countries. It also contributes significantly, even at subclinical levels, to morbidity and mortality from common childhood infections. Heightened awareness of the role of vitamin A in human health has led to an international effort to eliminate VAD and its consequences as a public health problem by the year 2000.

In 1987 WHO estimated that VAD was endemic in 39 countries based on the occurrence of clinical eye signs or symptoms, or very low blood levels of vitamin A ($<0.35 \mu\text{mol/l}$). This document updates this information based on biochemical evidence of subclinical VAD, i.e. population-based blood levels of vitamin A $\leq 0.70 \mu\text{mol/l}$, supported by other biological indicators and such ecological risk factors as diet. It is now estimated that VAD, including clinical and subclinical forms of severe and moderate degrees of public health significance, exists in 60 countries, and it is likely to be a problem in at least an additional 13 countries. An estimated 2.8 to 3 million preschool-age children are clinically affected, and 251 million more are severely or moderately subclinically deficient. At least 254 million children of preschool age are thus "at risk" in terms of their health and survival.

VAD is the result of two primary factors. First, there is a persistent inadequate intake of vitamin A to satisfy physiological needs. This is frequently exacerbated by secondary dietary circumstances such as insufficient consumption of dietary fat, which leads to inefficient absorption of this micronutrient. The second factor causing VAD is a high frequency of infections. Infections depress appetite, prompting an elevation in the body's vitamin A utilization and consequently the nutrient's inefficient conservation. Other conditions related to poverty, e.g. social underdevelopment (particularly affecting women); inadequate environmental sanitation; and insufficient water supply for growing food, drinking and maintaining adequate personal hygiene are associated with malnutrition generally, often including VAD. These conditions of deprivation are reflected in high infant and child mortality rates, which may be reduced to a degree by improved vitamin A status. It is important, therefore, to identify populations with subclinical VAD, as well as those with xerophthalmia, to quantify the at-risk population and to implement intervention programmes that reflect the true magnitude and severity of the problem. Recent methodological developments and refinements in their interpretation permit this to be accomplished far more readily than in the past.

This document provides a comprehensive listing of data on the magnitude and distribution of VAD based on clinical and biochemical parameters that are supported by ecological risk factors. The intention is to provide an overview of the global distribution of VAD. However, in the absence of, in particular, nationally representative data from all countries concerned, national prevalence estimates have been made based on the best available representative sub-regional data. Estimates of the numbers of people "affected" and "at risk" are thus approximations subject to refinement as more representative and comprehensive data become available. Nevertheless, it is clear that there is a significant VAD problem in at least parts of most countries in Africa, South and South-East Asia, and some areas of Latin America and the Western Pacific. Generally speaking, VAD is not believed to be of public health significance in countries with established market economies; however, most of these countries have not recently conducted surveys that would detect subclinical deficiency if it were to occur among their less affluent populations. Thus far little information is available from the former socialist countries of central Europe.

This document is a first step in providing updated national estimates of VAD and in generating regional and global figures on this basis. The data serve as a baseline for tracking progress nationally and globally towards the virtual elimination of VAD, which is one of the end-of-decade micronutrient goals endorsed by the World Summit for Children (1990), the International Conference on Nutrition (1992), and the World Health Assembly (1993).

The document is divided into four sections. The first section describes the nature of VAD and reviews the epidemiological issues involved in measuring and interpreting VAD prevalence studies. It also describes in some detail the steps taken in extracting data from various sources. The second section presents summary tables of the most recent VAD prevalence data by country and WHO region. The third section presents more detailed sub-national prevalence data by WHO region and, where available, sub-national areas, thereby showing the geographic variability of VAD within countries. Biological data are presented with information concerning the specific laboratory procedures employed. The fourth section provides complete bibliographic information for all data sources presented in the national and sub-national prevalence tables.

INTRODUCTION & BACKGROUND

VITAMIN A DEFICIENCY (VAD)

Vitamin A is an essential nutrient needed in small amounts for the normal functioning of the visual system, growth and development, maintenance of epithelial cellular integrity, immune function, and reproduction. VAD occurs when body stores are depleted to the extent that physiological functions are impaired even though clinical eye signs may not be evident. Because the vitamin is fat-soluble it is stored in the body when intake is in excess of physiological need. Nearly 90% of that which is stored is found in the liver. Depletion of stored vitamin A occurs over time when the diet contains too little to replace the amount used by tissues or reduced by breast-feeding. The level of depletion at which physiological functions begin to be impaired is not entirely clear. What is known, however, is that the vitamin is actively recycled through the liver and among tissues, and it appears that rates of utilization by specific tissues can at least partially adapt to diminishing availability. This adaptation and recycling serves to maintain relatively constant blood levels until body stores become depleted below a critical point for which adaptation can no longer compensate (1).

The integrity of epithelial barriers and the immune system are compromised before the visual system is impaired. This leads to increased severity of some infections and risk of death, especially among children (2). When vitamin A depletion is sufficient to affect the visual system, nightblindness occurs first due to the body's decreased ability to generate rhodopsin, which is essential for vision in dim light. This is accompanied by a loss of goblet cells from the epithelial tissue of the eye and results in xerophthalmia ("dry eye") which can affect both the conjunctiva (conjunctival xerosis and Bitot's spot) and cornea (corneal xerosis), and may lead to corneal ulceration, invasion by microorganisms,

and irreversible partial or total blindness (keratomalacia) (3).

Ocular symptoms and signs resulting from VAD-xerophthalmia have a long, well-recognized history, and have until recently been the basis for estimating the global burden from the disease. However, it is now recognized that the health of far larger numbers of preschool-age children (4), and perhaps older children and pregnant or lactating women as well, is compromised by VAD, even at moderate, and possibly mild, sub-clinical levels.

Figure 1 illustrates the relation between age and basal requirements for vitamin A on the basis of body weight and recommended safe levels of intake. Required intake during pregnancy and lactation are included in Table 1. The estimated





Fig. 1 Estimated basal requirements for vitamin A by age

basal requirements and safe level of intake for vitamin A by age group and sex are also given in Table 1. When weighted by a typical age-sex

distribution in a population, the basal vitamin A requirement for planning purposes is 250 μg daily and the recommended safe intake level is 550 μg .

Table 1. Estimated requirements for vitamin A (μg RE/day) (5)

Group	Age (years)	Basal requirement		Safe intake level	
		$\mu\text{g}/\text{day}$	$\mu\text{g}/\text{kg}$	$\mu\text{g}/\text{day}$	$\mu\text{g}/\text{kg}$
Both sexes	0-1	180	40-20	350	78-39
	1-6	200	13	400	26
	6-10	250	10	400	16
	10-12	300	8.3	500	14
	12-15	350	7.8	600	12
Girls	15-18	330	6.1	500	9.3
Women	18+	270	4.8	500	9.3
Pregnant women		370		600	
Lactating women		450		850	
Boys	15-18	400	6.1	600	9.3
Men	18+	300	4.8	600	9.3



*Vitamin A deficiency before, and 4 days after, receiving a single high-dose (200 000 IU) of vitamin A.
Photo: B. Underwood*

A 1989 estimation of the global supply of vitamin A is summarized in Table 2 according to WHO and UNICEF regions (6). In recent years a slight increase in total supply has occurred in all regions of the world except in parts of Africa where supplies, apart from North Africa, have remained unchanged or declined. The increase has occurred mainly in vegetable rather than animal sources of vitamin A with the exception of China where per capita intakes from animal sources have increased. Although the available global vitamin A per capita is above basal requirements, safe levels of intake are not met in South and South-East Asia, or in East and Southern Africa.

Per capita availability data must be viewed with caution. For example, the highest per capita availability of vitamin A-activity in foods is found in the Sahelian belt and West Africa, most of which is from vegetable sources of provitamin A. In these African regions, clinical VAD is endemic just as it is in South Asia where the available per capita supply, also largely from vegetable sources, is two-fifths as great. Clearly per capita data can



*Some vegetable sources of vitamin A.
Photo: B. Underwood*

Table 2. Available supply of vitamin A by WHO region ($\mu\text{g RE/day}$)^a

Region	Total	Animal	Vegetable
AFR	776	122	654
AMR	814	295	519
SEAR	431	53	378
EUR	738	271	467
EMR	936	345	591
WPR	997	216	781

Available supply of vitamin A by UNICEF region ($\mu\text{g RE/day}$)^a

Region	Total	Animal	Vegetable
Eastern & Southern Africa	453	137	316
Central & West Africa	1035	105	926
Middle East & North Africa	951	336	615
East Asia & the Pacific	850	168	682
Americas & the Caribbean	814	295	519
Devl./Indus. countries	NA	NA	NA
South Asia	435	71	364

^a Source: ACC/SCN

Table 3. Type of data available by WHO region

Region	Number of countries	No data or data prior to 1980	Data since 1980	National sample	Sub-national sample
AFR	46	19	27	7	20
AMR	36	22	14	1	13
SEAR	11	3	8	4	4
EUR	50	47	3	0	3
EMR	22	13	9	2	7
WPR	26	14	12	4	8

mask problems of uneven distribution. Hence, VAD continues to be a widely spread problem among preschool-age children even if recent information suggests that a decline is occurring in its most serious clinical form, keratomalacia (7).

Country data presented in this document are not all based on representative national sampling. A listing of the type of data available by region and used to derive the summary tables is given in Table 3. A country-by-country evaluation of the available data is provided in the notes for each country to assist readers in interpreting the prevalence data. In several countries, current information is not available and judgements were made based on projections as to how recent events would likely influence earlier prevalence data. Attention is drawn to the fact that data are not available or current for a large number of countries in those regions where VAD is widespread. Thus, the estimates in Table 4 reflect only available data and should not be interpreted as the total population "at risk" either regionally or globally. Based on the available data projected to reflect conditions in 1994, the global estimates of the numbers of children 0-4 years of age clinically affected is 2.8 million, and severely and moderately subclinically affected (taken as one category), is 251 million. These estimates are provided by country in the summary tables and are summarized by WHO region in Table 4.

Table 4. Estimates of affected and at-risk populations*

Region	Clinical (x10 ⁶)	Subclinical severe & moderate (x10 ⁶)	Prevalence (%)
AFR	1.04	52	49.0
AMR	0.06	16	20.0
SEAR	1.45	125	69.0
EUR	NA	NA	NA
EMR	0.12	16	22.0
WPR	0.13	42	27.0
Sub-Totals	2.80	251	
TOTAL		254	

*Based only on a projection for 1994 from those countries in each region where data were available. See Table 3 for numbers of countries without data.

EPIDEMIOLOGY OF VAD

VAD as a public health problem occurs within an ambience of ecological, economical and social deprivations in the macro-environment in which populations are found (i.e. regions and countries), and in the micro-environments in which families live (i.e. communities and households). The rela-

tive influence of causal factors at both the macro- and micro-level will vary among countries, and even regions within countries, necessitating a situational analysis to understand and subsequently design appropriate and effective intervention programmes to change specific undesirable situations. It is, therefore, with considerable reservation that national and global projections are made that could be inappropriately applied to local situations. Nevertheless, there are some underlying epidemiological traits that tend to characterize most situations where VAD occurs as a public health problem (8). These are described below.

Ecological factors

At the macro-level, hostile environments, e.g. arid, infertile land, or the periodicity of excessive rain and humidity, in part determine the variety and amount of foods rich in vitamin A-activity that can be grown, and the duration of their availability. This applies particularly to vegetables (e.g. green leafy vegetables), and fruits that require abundant water supplies and/or moderate temperature to grow. Where the necessary favourable growing conditions occur in food-scarce countries, even if only seasonally, national agricultural policies generally favour production of staples as food for local populations. From a national perspective, vegetable and fruit crops are of less importance and thus do not compete for land use. Crops rich in vitamin A-activity for local consumption, therefore, are more often provided through horticultural activities at the micro-level. Even at community and household levels, however, the characteristics of a hostile environment, particularly where water is in short supply, limit home and community gardening activities and, as a consequence, the availability of inexpensive sources of vitamin A. Thus, countries or parts of countries with long periods of water shortage and relatively constant hot temperatures are more likely to have a VAD problem than those with stable water supplies.

The seasonality of VAD is only partially related to ecologic factors that influence food availability. The pattern of disease frequency is also important. VAD tends to reach its apex following the peak prevalence of diarrhoeal and respiratory diseases. Overcrowded housing and contaminated environments associated with poor living conditions contribute to the problem. Measles epidemics that occur under these conditions are especially devastating and often precipitate VAD, frequently resulting in blindness and death for many children.

Social factors

Social underdevelopment within a country limits accessibility to health and social services, including education. Under-educated, impoverished women tend to follow traditional ideas and practices, and are less confident in engaging in social interactions where more modern concepts and practices are promoted. Due to under-education, they are less likely to learn from educational materials typically displayed at health centres and used in health-related community educational activities, including those concerned with appropriate child care and feeding practices. Under-educated males also are less likely to adopt within their households new ideas and practices related to family care and feeding. A socially backward, impoverished environment also favours large families with consequent overcrowding that is associated with poor environmental sanitation and personal hygiene. As noted above, these are prime conditioning factors for VAD and malnutrition.

Economic factors

Poverty is a root, though not invariable, cause of VAD in public health terms. Because only foods of animal origin contain preformed sources of vitamin A, which are generally relatively expensive, VAD is confined largely to impoverished

countries, neighbourhoods and families which rely on less expensive provitamin A sources to meet their requirements. Provitamin A sources must be converted to retinol before they can provide protection from VAD. The series of events between consumption of provitamin A and its conversion to retinol include several steps that are dependent on normal physiological functions. For reasons discussed below under *Host factors*, it is more difficult to satisfy vitamin A-activity needs of infants and young children from foods of vegetable origin than from other food sources.

Poverty contributes in other ways, some already noted, to inadequate living conditions that are associated with high death rates among infants and young children. Unemployment and low-wage jobs are major obstacles to overcoming VAD in depressed environments.

Clustering

As already mentioned, the occurrence of clinical VAD tends to cluster rather than to be evenly distributed. Clustering within countries at the macro-level is related to ecological factors noted above exacerbated by poorly developed infrastructures to distribute vitamin A-containing foods from excess-to deficient-areas. Because vitamin A-rich foods tend to be quite perishable, they are especially susceptible to inadequate intra-country distribution.

Clustering has been described primarily based on the occurrence of clinical eye signs. It is likely to reflect the convergence of several risk factors that lead to depletion of vitamin A stores in the surrounding child population among which a few individuals who have been exposed to additional causal factors have developed clinical deficiency. For this reason severely subclinically deficient populations of children up to 5 years of age, based on the distribution of serum retinol levels, are

considered to be as much at risk of severe morbidity and mortality as those populations experiencing clinical deficiency. Moderately subclinically affected populations are also likely to be at higher risk but the magnitude of risk is unknown.

Host factors


Age. Varying levels of VAD can occur at any age, from subclinical effects that increase risk of morbidity and mortality to blinding malnutrition (keratomalacia). As a public health problem, however, VAD affects children of preschool age because of their great susceptibility to infections and due to an increased demand for the micronutrient by the body to support their rapid growth. The potentially blinding corneal disease is most prevalent among children under 3 years of age and is usually associated with PEM. An increased risk of death of at least 60% is associated with severe, potentially blinding VAD malnutrition (9). The mortality risk associated with VAD of lesser severity extends at least from 6 months to 5 years of age, and perhaps beyond. The elevated risk of death among those less severely clinically affected, and severe to moderately subclinically affected, is estimated to be about 23% (4).

There is little information regarding the health consequences of VAD among school-age children. The prevalence of mild xerophthalmia, notably Bitot's spots, may be highest in the school-age group, although this may be more a reflection of past rather than current vitamin A status (10).

Sex. No consistent sex difference in vulnerability is demonstrated based on physiological parameters. Differences have been reported from some cultures, which are more likely to be related to sex differences in cultural practices of feeding and care rather than to physiological differences.

Feeding practices. Breast milk provides retinol in a readily absorbable form. Clinically apparent VAD is rare among populations where breast-feeding prevails. Even though clinical deficiency rarely occurs as long as a child is receiving breast milk, depletion of an infant's body stores, leading to subclinical deficiency and consequent health risks, may occur by six months of age when maternal vitamin A status is inadequate and thus, breast milk vitamin A content is low (11). In general, the problem of subclinical depletion increases in significance between 6 months and 3 years of age during which complementary foods and later the family diet, represents a large proportion of the infants diet. These foods often do not contain vitamin A in amounts that adequately replace that provided from the diminishing contribution of breast milk. The diet of the newly weaned child frequently has very little vitamin A and often contains less fat than at any other period in the life cycle. Dietary fat is especially important for the absorption of vegetable sources of provitamin A. The post-weaning period, until a child has begun receiving a diversified family diet, is therefore one of great vulnerability to VAD.

Disease patterns. The frequency, duration and severity of infections contribute directly and indirectly to vulnerability. Infections influence appetite and are especially devastating for the weaned child. Infections lessen efficiency of absorption, conservation and utilization of vitamin A. Frequent acute bacterial infections damage mucosal surfaces required for absorption. Furthermore, intestinal worm infections may directly compete for uptake of vitamin A in addition to their more general impact on health by suppressing appetite. The frequency of diarrhoeal and respiratory infections is associated with VAD vulnerability (12). For diarrhoeal disease, restoring vitamin A status decreases the severity of subsequent episodes and the risk of death (13). Curiously, no such link has yet been established with respiratory illness, except for pneumonia associated with measles (14).



*Breast-feeding reduces the risk of VAD.
Photo: B. Underwood*

Periods of increased physiological need. Vitamin A needs on a body-weight basis are increased during periods of rapid growth (see Table 1 and Fig. 1), which is one reason for the greater vulnerability of younger children. School-age children are growing but at a slower rate than at earlier ages, at least until adolescence. In areas where VAD is endemic, however, the prevalence of Bitot's spots in schoolchildren is often above that seen in younger age groups, and may not in




Fig. 3 The trap of deprivation

fact be responsive to improved vitamin A status (15). There is only limited information to determine if increased morbidity or mortality risks are associated with Bitot's spots in school-age children.

Other periods of increased physiological need are gestation and lactation. Women are vulnerable to VAD in these periods because of the increased need to provide for the developing fetus and, following parturition, to replace vitamin A transferred from maternal stores via breast milk to the nursing infant. Pregnant and lactating women in underprivileged populations often report night blindness and studies have found that their breast milk frequently is low in vitamin A (breast milk vitamin A content reflects that of maternal diet and maternal vitamin A status) (11,16). Few data are available to determine if there is an increase in morbidity or mortality risk to the mother that is associated with mild depletion of vitamin A stores. For this reason, pregnant and lactating women are not included in the global estimates of the at-risk population. This will be corrected as the MDIS databank receives more specific information about the vitamin A status of pregnant and lactating women, e.g. data on breast milk vitamin A levels.

INDICATORS OF VAD

Clinical eye signs are rare events. Therefore, when examinations are performed to determine prevalence of severe, potentially blinding VAD, large representative samples are required. Eye signs are, however, the most specific and sensitive of the VAD indicators. Biological indicators of subclinical VAD below selected cut-off points are more common, thus requiring a smaller sample size for estimating prevalence. Regrettably, however, all the subclinical indicators available to date lack specificity and/or sensitivity. Furthermore, when these indicators are used as the basis for cut-off points in determining the severity

of vitamin A deficiency, additional population-based validation is required.

Clinical indicators

There is a standard classification system for ocular indicators of VAD, and minimum prevalence criteria for interpretation to identify a public health problem have been widely accepted and applied (Table 5). These criteria have been used to identify countries in the database with a significant public health problem of xerophthalmia and hence risk of VAD-related blindness.

Serum retinol

Distribution curves for serum or plasma retinol levels are useful for identifying populations likely to be at risk of VAD. The prevalence of values below selected cut-off points is useful for estimating the relative risk and prevalence of severity and its magnitude as a subclinical problem. A prevalence of values $\leq 0.35 \mu\text{mol/l}$ above 5% has been used globally to define a deficient population as corroborative evidence of clinical eye signs for risk of blinding malnutrition. However, this cut-off is insufficient to identify the prevalence of risk to those likely to have inadequate vitamin A status, i.e. who are subclinically deficient and suffering health consequences. For this purpose, $<0.70 \mu\text{mol/l}$ was suggested earlier as a cut-off for inadequate vitamin A status, but the prevalence criteria to define a public health problem was not generally applied except in the Latin America and Caribbean region (AMR). In the Region of the Americas a prevalence of $>15\%$ below $0.70 \mu\text{mol/l}$ has been generally applied to identify an unacceptable situation (18). New evidence substantiating increased mortality risk among populations without clinical signs has necessitated a reevaluation of cut-off points and criteria for interpreting serum retinol distributions. A WHO consultation (19) held to address this question recommend that

Table 5. Classification of xerophthalmia and prevalence criteria constituting a public health problem (17)

Criteria*	Minimum prevalence
Night blindness (XN)	>1.0%
Bitot's spot (X1B)	>0.5%
Corneal xerosis and/or ulceration (X2,X3A,X3B)	>0.01%
Xerophthalmia-related corneal scars (XS)	>0.05%

(A serum retinol level $<0.35\mu\text{mol/l}$ in $>5\%$ of the population is corroborative evidence)

* See reference 3 for detailed definitions and illustrations of the clinical classification and diagnosis of xerophthalmia.

a cut-off level of $\leq 0.70\mu\text{mol/l}$ be used and that the prevalence of values below the cut-off be ranked to indicate the degree of public health importance as shown in Table 6. The consultation recommended that a severe public health problem be defined as existing when 20% or more of a surveyed preschool-age child population has values $\leq 0.70\mu\text{mol/l}$, and a minimum prevalence of $\geq 10\%$ to identify a moderately severe public health

problem. It was also recognized that among well-nourished, healthy populations of preschool-age children, and even those still living in poverty but whose vitamin A status is adequate, fewer than 5% have values $\leq 0.70\mu\text{mol/l}$ (20). Therefore, a category is included to identify a mild problem worthy of consideration even when the prevalence is $<10\%$ of the surveyed preschool-age population having serum levels $\leq 0.70\mu\text{mol/l}$.



Unilateral blindness before 6 months of age due to vitamin A deficiency in a non-breast-fed Thai child given an unfortified condensed milk product. Photo: E. Wasanwisut

Table 6. Prevalence of VAD in children ≥ 1 year of age of serum values $\leq 0.70 \mu\text{mol/l}$

Level of public health problem	Prevalence
Mild	≥ 2 - $<10\%$
Moderate	≥ 10 - $<20\%$
Severe	$\geq 20\%$

The prevalences in Table 6 for severe and moderate subclinical VAD have been used in this document to classify countries and to estimate the at-risk population. The "mild" category was not included in calculating the at-risk estimates. These prevalences need further verification under field conditions using other indicators of VAD. Several factors common to deprived populations can lower serum values of vitamin A or status independently of intake, e.g. acute and chronic infections (21). The variable prevalence of these confounders among populations may in future necessitate using different criteria for different situations.



Young girl with xerophthalmia consuming a low-vitamin A meal. Photo: International Center for Eye Health, London

Other biological indicators

As noted above, additional supporting evidence is needed because no single biological indicator of subclinical VAD by itself, is of sufficient specificity and sensitivity to identify a public health problem. Other indicators of subclinical VAD include dose response tests and CIC. These measures of VAD also have limitations both due to confounders and in terms of reliability of interpretation. In addition, they have had very limited application in field surveys. Data using these indicators are entered into the database as they accumulate. However, data collected on the basis of these new assessment techniques, have not been included here for determining prevalence of at-risk populations. They are used only as corroborative information. The prevalence of VAD can be determined only by using biological indicators. Once again, the root cause of VAD in a public health context is habitual inadequate dietary intake of foods containing vitamin A-activity, i.e. pre-formed vitamin A and provitamin carotenoids. Reliable quantitative assessment of habitual dietary intake is problematic, however. Newer approaches to solving this problem use qualitative or semi-quantitative measures of intake frequency of vitamin A-containing foods. The information obtained is useful in corroborating biological indicators, which is how it has been used in this document.

A fuller discussion of the use of other nutritional, ecological and demographic indicators for surveillance purposes other than determining prevalence is presented in a separate WHO document.¹

¹ *Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes document, WHO/NUT/94.1, available from the Nutrition unit, WHO, Geneva.*

REFERENCES

1. Blomhoff R. et al. Vitamin A metabolism: New perspectives on absorption, transport, and storage. *Physiological reviews*, 1991, **71**:951-990.
2. Ross CA, Hämmerling UG. Retinoids and the immune system. In: Sporn MB, Roberts MB, Goodman DS, eds. *The Retinoids: Biology, Chemistry and Medicine, 2nd edition*. New York, Raven Press, Ltd. 1994:521-543.
3. Sommer A. *Vitamin A deficiency and its consequences: A field guide to their detection and control, 3rd ed.* Geneva, World Health Organization, 1995.
4. Beaton GH et al. *Effectiveness of vitamin A supplementation in the control of young child morbidity and mortality in developing countries*. New York, United Nations Administrative Committee on Coordination/Subcommittee on Nutrition, 1993 (ACC/SCN State-of-the-Art Series, Nutrition Policy Discussion Paper No. 13).
5. *Requirements of vitamin A, iron, folate and vitamin B₁₂. Report of a joint FAO/WHO Expert Consultation*. Rome, 1988 (Food and Nutrition Series, No. 23).
6. ACC/SCN. *Second report on the world nutrition situation, Vol.1. Global and regional results, October, 1992*. Vol. II, March 1993.
7. ACC/SCN. *Controlling vitamin A deficiency*. ACC/SCN State-of-the-Art Series, January 1994 (Nutrition Policy Discussion Paper No. 14).
8. Underwood BA. The epidemiology of vitamin A deficiency and depletion (hypovitaminosis A) as a public health problem. In: Livrea MA, Packer L, eds. *Retinoids: Progress in research and clinical applications*. New York, Marcel Dekker, Inc., 1993:171-184.
9. *Prevention of childhood blindness*. Geneva, World Health Organization, 1992.
10. Sommer A, Emran E, Tjakrasudjatma S. Clinical characteristics of vitamin A responsive and nonresponsive Bitot's spots. *American journal of ophthalmology*, 1980, **90**:160-171.
11. Underwood BA. Maternal vitamin A status and its importance in infancy and early childhood. *American journal of clinical nutrition*, 1994, **54** (Suppl):5175-5245.
12. Sommer A, Tarwotjo I, Katz J. Increased risk of xerophthalmia following diarrhea and respiratory disease. *American journal of clinical nutrition*, 1987, **45**:977-980.
13. Barreto ML et al. Effect of vitamin A supplementation on diarrhoea and acute lower-respiratory infections in young children in Brazil. *Lancet*, 1994, **344**:228-231.

14. Coutsoudis A, Broughton M, Coovadia HM. Vitamin A supplementation reduces measles morbidity in young African children: A randomized, placebo-controlled, double-blind trial. *American journal of clinical nutrition*, 1991, **54**:890-895.
15. Semba RD et al. Response of Bitot's spots in preschool children to vitamin A treatment. *American journal of ophthalmology*, 1990, **110**:416-420.
16. Newman V. Vitamin A and breastfeeding: A comparison of data from developed and developing countries. Wellstart, 1993.
17. *Control of vitamin A deficiency and xerophthalmia. Report of a joint WHO/UNICEF/USAID/Helen Keller International/IVACG meeting.* Geneva, World Health Organization, 1982 (WHO Technical Report Series, No. 672).
18. *Manual for nutrition surveys. Second edition. Interdepartmental Committee on Nutrition for National Defense.* Maryland, National Institutes of Health, 1963.
19. *Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. Report of a joint WHO/UNICEF consultation.* Geneva, World Health Organization, May 1994 (Review version, WHO/NUT/94.1).
20. Flores H et al. Serum vitamin A distribution curve for children aged 2-6 y known to have adequate vitamin A status: a reference population. *American journal of clinical nutrition*, 1991, **54**:707-711.
21. Filteau SM et al. Influence of morbidity on serum retinol of children in a community-based study in northern Ghana. *American journal of clinical nutrition*, 1993, **58**:192-197.

MDIS

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY

TABLES SUMMARIZING VITAMIN A DEFICIENCY PREVALENCE BY COUNTRY IN EACH WHO REGION

INTERPRETING DATA IN THE MDIS

To ensure their proper interpretation there are several considerations which must be borne in mind in the present analysis and presentation of VAD prevalence data. Although efforts have been made to standardize methodologies employed in the design and implementation of VAD prevalence surveys, there is in fact considerable variation in the way these surveys were conducted and how results were analysed. This means lack of comparability across studies.

Survey design

Depending upon the objectives for VAD surveillance, distinctly different approaches have been used in the design of surveys and the selection of samples to ensure the greatest degree of representativeness. For cross-sectional surveys, when data are collected at a single point, it is generally appropriate to use stratified sampling techniques with sample sites being selected to provide a representative impression of a population, using either random, cluster or PPS (population proportional to size) techniques. Since clinical VAD is rare, even in areas where there is a significant public health problem, i.e. 1 case of corneal xerophthalmia seen in 1000 children ($X^2 = .01\%$) the sample size requirements for clinical surveys must be quite substantial to ensure reliability. It is crucial, therefore, that sample size calculations be considered when reviewing data and when ascertaining the precision of individual survey estimates. Furthermore, since in most developing countries, VAD tends to occur only in a small number of communities (clustered) rather than being equally distributed throughout populations (homogenous) specific survey sampling designs are required so that variance estimates are not miscalculated and confidence intervals misrepresented.

A survey methodology influences the interpretation of results in terms of their representativeness and statistical precision, and different methodologies make inter-country

comparisons difficult. The MDIS has attempted to deal with this problem by cataloguing the sampling methods used in each survey, including the design, selection of sites, selection of individuals, and the sample size. However, many documents which summarize information from VAD prevalence surveys do not provide complete information on the study methodology.

Subject selection

The MDIS has also included information on subject characteristics: sex, age, and sub-national residence. This information is particularly important for comparing prevalence estimates across studies. There are important differences in the prevalence of VAD in various age groups, and so direct comparisons are difficult. It is imperative to detail this information when presenting VAD prevalence data.

Data aggregation

For purposes of global advocacy, an understanding of the national magnitude of VAD is required. However, the aggregation of data to derive a single national estimate limits the ability to highlight the important differences that may exist in the distribution of VAD within countries. In the MDIS, data are maintained for sub-national areas, so that the intra-country disparities in VAD may be seen, and populations at greatest risk recognized.

Limitations

The data presented in this document are the most up-to-date vitamin A prevalence information available to the MDIS based on an extensive literature review and reports available to WHO and other organizations. These estimates will be revised periodically as more data become available. It is recognized that some reports may have been missed because they were not published,

or otherwise unavailable to WHO. One of the prime objectives of this document is to identify and fill remaining gaps in the global database, and to gain access to supplementary information that will ensure that subsequent revisions are complete.

Surveys from small areas may provide a biased prevalence estimate, especially if they are performed in areas known to have a high prevalence of VAD, and may not represent the entire country. The present document makes no claims about the accuracy of laboratory procedures performed in the surveys, the comparability of the cross-survey assessments, or the methods employed in each survey. For more details about the characteristics of individual surveys, the reader is directed to original documents as specified in the *Bibliographic references* section. These references are on file as part of the MDIS database in the WHO Nutrition unit.

GUIDE TO USING SUMMARY TABLES

Each country where a problem of VAD is documented or suspected is listed by WHO region. Where data are based on surveys prior to 1980, judgement is used in determining how recent developments may have altered the situation. An "X" designates the estimated category of deficiency, i.e. clinical, severe subclinical or moderate subclinical where survey data are not available but current reports indicate there is a problem. When a rate is derived from more than one survey, a superscript E is inserted by the number. Where known, the year of the survey on which the prevalence is based is given. The national population figure is the UN projection for 1995 of the 0-4-year-age group. Prevalence values in the table are therefore for preschool-age children only.

Criteria used to define the severity of a public health problem are as follows:

Clinical. The number in the table refers to the prevalence of total xerophthalmia, or of other

clinical eye signs, i.e. Bitot's spot (X1B), corneal xerosis (X2), keratomalacia (X3) and corneal scars (XS) and/or symptoms, i.e. nightblindness (XN) measured. Values are included only if they exceed the level defined by WHO as constituting a public health problem as defined in Table 5. Where clinical data are documented the population clinically affected is noted (in thousands) in parentheses directly below the prevalence (%).

Severe subclinical VAD. A prevalence of $\geq 20\%$ with blood values $\leq 0.70 \mu\text{mol/l}$ (with or without clinical eye signs or symptoms) as shown in Table 6.

Moderate subclinical. A prevalence of ≥ 10 - $<20\%$ with blood values $\leq 0.70 \mu\text{mol/l}$ (with or without clinical eye signs or symptoms) as shown in Table 6.

A multiplication factor was derived for countries where representative national surveys of vitamin A deficiency were unavailable. Where sub-national surveys were available, extrapolations were made to the proportion of the total country likely to be affected considering similar ecological conditions. From this a multiplication factor was generated as shown in Table 7:

Table 7. Method for determining at-risk population


Estimated portion of country affected	Multiplication factor
National sample	0.75
>60-<75%	0.60
>30-<60%	0.40
>20-<30%	0.25

The at-risk population was estimated as follows. The multiplication factor was applied in countries in which WHO criteria to identify a public health

problem of VAD clinically or subclinically were met. The total population of children 0-4 years of age according to the UN projection for 1995 multiplied by the factor determined the total, i.e. clinically + subclinically, at-risk population. In countries with clinical VAD, the prevalence of clinical signs was multiplied by the total at-risk population to estimate the number of children clinically affected. That number is given in parenthesis under the clinical prevalence figure. In countries with no clinical problem but a severe subclinical one, the at-risk population estimate includes those who also are moderately subclinically deficient.

Countries categorized by degree of public health importance of vitamin A deficiency

April 1995




The designations employed and the presentation of material on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines represent approximate border lines for which there may not yet be full agreement.

Prevalence of Vitamin A Deficiency

African Region

April 1995



WHO 95216

The designations do not necessarily imply VAD severity is uniformly distributed throughout each country

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Algeria		4260				?	No data available. VAD likely.	
Angola	1973	2195	3.0 (26.3)		X	0.40	873	No recent data available. Survey in Bie Province (1973) found a X1B prevalence of 3%. VAD currently suspected to be a problem as a result of malnutrition brought on by war, drought and massive migration. Further assessment needed.
Benin	1989	1061	8.3% (52.8)			0.60	636	No national data. VAD considered a problem in clustered areas in the northern sections of the country. A study in North Atacora (1989) found a XN rate of 3.4% and a X1B rate of 4.9%. Older data available on an unidentified age group in 3 districts of Atacora (1975) found a high xerophthalmia rate of 3.35%. No biochemical data are available, though clinical data suggest that VAD is a public health problem. Vitamin A capsules are being distributed by UNICEF. Further assessment needed.
Botswana	1992	238			X	0.25	59	ICN report (1992) shows evidence of mild-to-moderate vitamin A deficiency. A nutritional survey (1971) of 77 255 people in 3 villages recorded very few clinical signs of deficiency.
Burkina Faso	1986-89 1978	1899	1.7-3.3 (28.2)	70.5		0.60	1139	No national data. Sub-national survey (1986) of three Northern provinces of the country (Yatenga, Passore and Sourou) found a X1B rate of 0.27%, a XN of 2.82% and a total xerophthalmia rate of 3.27% in children 0-5 years of age. A survey of six provinces (1989) found a XN rate of 1.42% and a X1B rate of 0.26% among children aged 2-10 years. A dietary survey (1984) found low intakes of fruits while an older survey (1978) observed high level of marginal VAD, with 70.5% of preschool children having serum retinol levels <0.70 µmol/l and an overall mean of only 0.58(± 0.23) µmol/l. HKI, UNICEF and MOH distributed vitamin A capsules in four northern provinces between 1986 and 1989.
Burundi		1183		X		0.40	473	No data available. VAD highly likely in parts of country due to recent civil unrest and occurrence in surrounding countries.

* Refer to *Interpreting data in the MiDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Cameroon	1992	2270	0.5 (4.5)	19.7		0.40	908	No national data. Study in the Extreme North Province (1992) found a X1B prevalence of 0.47%, the highest prevalence found in the mountainous region and that 19.7% of children in periurban and flooded areas had serum retinol levels <0.70 µmol/l. A small study in Kale Department of the North Province (1988) found 50% of 3-12-month-old children with serum retinol levels <0.35 µmol/l.
Cape Verde		67		X		0.40	27	Though there are no data on children under 5 years of age, data from surveys of the general population in Fogo and San Vicente suggest high prevalence of VAD with X1B rates of 2.9% in females and 3.6% in males. However, these data are based on very small sample sizes and their precision is questionable. Biochemical data from the same study of children between 7-13 years of age noted 6.7% with serum retinol levels <0.35 µmol/l.
Central African Republic		621					?	No data available. VAD likely.
Chad	1986	1108	2.7-4.5 (29.9)			0.75	831	No national data. Data on 0-5-year-olds from Chari-Baguirme, Bath and Ouaddi (1986) in the Northern and Central areas of the country found clinical VAD levels extremely high with a total xerophthalmia rate of 2.7%. Though there is considerable variation between regions (rural areas of Chari-Baguirme reporting a rate of 4.5%), VAD is considered a public health problem in the country. No biochemical data are available.
Comoros		130					?	No data available.
Congo	1988	483		26.0*		0.25	121	No national data. Biochemical data collected from Brazzaville and Pointe Noire (1988) found low serum retinol levels among children, especially those with malaria (83.3% <0.70 µmol/l). The mean was 0.52 µmol/l.
Côte d'Ivoire	1994	2927		46.6		0.60	1756	Preliminary study in 4 departments in the northwest (1994) found serum retinol levels of <0.70 µmol/l in 46.6% of the infants surveyed after supplementation. A survey in 3 communes (1988) found that 25% of children aged 1-3 years had serum retinol levels <0.35 µmol/l.
Equatorial Guinea		70					?	No data available. VAD likely.

* Refer to *Interpreting data in the MIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Eritrea						?	No data available. Study planned for 1995.	
Ethiopia	1980-81	9567	4.6 (330.1)	51.0		0.75	7175	National data (1980-81) from representative survey of preschool children aged 6 months-5 years indicated VAD to be a significant public health problem in all zones of the country except the Ensete zone, with a national X1B average of 1.0% and estimated total xerophthalmia rates of 6%. Large-scale study in Shoa region (1981) found X1B prevalence just above 0.5%. Biochemical data (1981) corroborated this finding with 59.6% of children 6-71 months of age having serum retinol levels <0.70 µmol/l and a mean of 0.62 µmol/l. VAD has also been found to be especially severe in refugee camps.
Gabon		231					?	No data available. VAD likely.
Gambia	1988	169		X		0.25	42	Data from surveys (1988) of children 2-6 years of age indicate that VAD is highly seasonal, occurring during the months of January-March and October-November. There was a large increase in mean serum retinol levels from 0.41 µmol/l in January to 0.78 µmol/l in August.
Ghana	1989-91	3049	1.1% (13.4)	54.9*		0.40	1226	Data from Kassena-Nankana (1989-91) indicate that VAD is a problem in the Northern and Upper East areas of the country. Over 14% of the preschool children who had serum samples taken had serum retinol levels <0.35 µmol/l and over 50% <0.70 µmol/l. A XN rate of 1.04% and a X1B rate of 0.01% was found. Morbidity Survey (1991) found an XN rate of 1.5% while the Survival Survey (1991) found 0.70%. Large-scale vitamin A supplementation trial has increased awareness of the VAD problem and has led to the establishment of a national VAD control program by MOH.
Guinea		1308					?	No data on children <5 years of age. Data on pregnant women (1980) show some problems of VAD with X1B rates as high as 5.2% in the north of the country. VAD is suspected also to affect children and the elderly.
Guinea-Bissau		179					?	No data available. VAD likely.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Kenya	1976-81	5162	0.2-1.1 (13.4)			0.40	2065	National data (1976-81) on an unspecified age group showed low prevalences of VAD (0.26%) while regions in the Highland tribes of Baringo and pockets of arid and semi-arid zones (Marsabit, Mwea) indicated VAD to be of public health significance with X1B rates as high as 1.13%. Hypovitaminosis A is thought to be a problem among a large proportion of children.
Lesotho	1993	295	0.0	78.0		0.75	221	A nationally representative survey (1993) found no clinical signs among children 2-3 years of age and only a few signs of Bitot's spots in children aged 4-6 years. More than 78% of preschool children had serum retinol values <0.70 µmol/l and 13.4% <0.35 µmol/l. A mean of 0.6 µmol/l was recorded.
Liberia		570					7	Nutritional assessment (1990) of Guinean nationals and Liberian refugee children in Guinea showed no clinical signs of avitaminosis A. Mild VAD likely.
Madagascar		2598						Small survey conducted in 1992. Data not available. Mild problem likely.
Malawi	1983-89	2323	2.6 [#] (27.9)			0.60	1394	Survey in the Lower Shire Valley (1983) revealed that VAD was of public health significance in this area with a total xerophthalmia rate of 3.0%. A study in Mbalachanda District (1988) found a prevalence of only 0.6% while a survey in Mkhota (1989) found a xerophthalmia rate of 1.7%. Old data analysis in the Lower Shire Valley (1970) found that 95% of the households had inadequate vitamin A intakes during the post rainy season. The health sector distributes vitamin A supplements to high-risk groups.

* Refer to *Interpreting data in the MIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Mali	1986 1978-79	2131	5.5 ^b (103.9)	73.0 ^b		0.75	1592	National data estimated from surveys done in most regions of the country. Data from 3 different populations in Tombouctou and Segou (1986) found all xerophthalmia rates to be above those indicative of posing a public health problem. Biochemical study (1990) conducted in NW Bamako noted that 15.1% of children 2-10 years of age had serum retinol levels <0.35 µmol/l. Biochemical data from Southern Mali (1978-79) indicated a high prevalence of subclinical VAD with over 12% of children 0-6 years of age having serum retinol values <0.35 µmol/l with a mean of 0.58(± 0.2) µmol/l. There are negligible differences in serum retinol levels between the rainy and dry seasons. In a survey of four populations (1986), each was found to have total xerophthalmia rates above 5% in children <5 years of age. A survey in Douen (1990) found a XN rate of 9.0%. In 1989 as part of a 5-year programme, vitamin A capsules were distributed to pregnant women and children under 5 years of age by HKI, USAID and AED.
Mauritania	1983 1987	428	2.6 ^b (6.7)	41.6 ^b		0.60	257	Study by USAID (1983) found xerophthalmia prevalences between 2.5% and 2.7%. More recent data from Nouakchott (1989-90) observed significantly lower clinical prevalence rates for children aged 7-10 years. Biochemical data from Hodh-El Gharbi (1987) showed a wide variation in serum retinol levels, with 2/3 of all children aged 1-15 years in Kerkerat having serum retinol values <0.70 µmol/l but only 10% of children in Limberha falling below this cut-off point.
Mauritius		1130					1	No data available. Vitamin A deficiency not considered a public health problem.
Mozambique	1990	2943	0.7 (5.2)			0.25	736	No national data. Survey in 3 cities (1990) of 10 267 children aged 6 months to 6 years found a VAD prevalence of 0.7%. No biochemical data are available. Clinical information also available from a survey of 3790 displaced children of war in Zambezia Province (1987-88) where it was noted that 11% of all childhood blindness was due to keratomalacia, with estimated prevalences of XN=7.3%, X3B=0.25% and XS=0.89%.

* Refer to *Interpreting data in the MIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Namibia	1992	304			20.4	0.75	228	Biochemical data available from survey conducted in 17 villages from 4 regions (1991) which revealed that 20.4% of children 2-6 years of age had serum retinol levels below $<0.70 \mu\text{mol/l}$ and 3.1% $<0.35 \mu\text{mol/l}$. However, there were quite dramatic differences in serum vitamin A levels between villages, with only 2 villages having any cases of severe subclinical VAD (serum retinol $<0.35 \mu\text{mol/l}$). It is noteworthy that the sample size for the survey was quite small.
Niger	1988	1820	2.0 (27.3)			0.75	1365	Survey data from Tahoua, Maradi and Zinder (1986) demonstrated a severe VAD problem with XN=4%, X1B=0.72% and corneal lesions of 14/1000 preschool children. National survey (1988) found a 2.0% prevalence rate for nightblindness. Data from several regions (1990) indicated that VAD is widely distributed, with XN >3% in Tera, Tillaberi and Ouallam Provinces. Dietary survey of 6-71-month-old children completed in 1994 in Tahoua and Maradi. Biochemical data (1970s) found the surveyed refugees to be severely vitamin A deficient.
Nigeria	1994	23 409	1.0 (375.6)			0.75	17 557	National survey (1994) found a XN rate of 1.0% among children <6 years of age. Biochemical data from the south found a serum mean of $0.72 \mu\text{mol/l}$. Study (1977) noted that 4% of blindness in children was due to keratomalacia and a later study in Kaduna Eye Hospital (1979) claimed that 69% of all blind children had corneal ulceration. Biochemical data in Cross River State (early 1980s) showed a mean serum level of $1.1 \mu\text{mol/l}$ among children aged 0-1 years.
Rwanda	1987	1720	1.3 (13.4)	X		0.60	1032	No national data. Sub-national data from Gikongoro Prefecture (1987) revealed a X1B rate of 1.3% among children 0-6 years of age. National survey was scheduled for 1993. Due to deteriorating conditions, VAD likely to be a country-wide problem.
Sao Tome & Principe							?	No data available.

* Refer to Interpreting data in the MIDS pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Senegal	1988 1991	1454	0.0	71.5		0.60	872	No national data. Study in Malicounda (1990-91) found no signs of xerophthalmia in children 2-14 years of age. Survey in Gadiack and Diop N'doffene (1989) in April and June before and after the mango season found 40% and 11.4% <0.35 µmol/l, respectively. Data from Diourbel and Fatick (1988) indicated that 14.1% of the 1-6-year-olds presented signs of XN, although only marginal X1B rates noted. Biochemical results on children normal by ICT showed a mean of 0.45 µmol/l. Survey in Linguere (1991) showed a mean of 0.61 µmol/l (71.5% <0.70 µmol/l) and that 90% of the diets sampled contained less than 50% of the RDA of vitamin A.
Seychelles							?	No data available. No problem likely.
Sierra Leone		876			X		?	Vitamin A deficiency has not been reported in Sierra Leone. This may be due to the fact that oils (palm oil in particular) are widely consumed. However, it is likely that there is a moderate subclinical problem.
South Africa	1991	5838	0.0	49.0		0.25	1459	Survey done in Bester Farm (1991) found that 49% of those surveyed had serum retinol levels <0.70 µmol/l and 5% had levels <0.35 µmol/l. No clinical signs of VAD were found.
Swaziland		136					?	No data available. VAD likely.
Togo	1992	764	10.0 (45.8)			0.60	458	National survey (1992) indicates vitamin A is endemic in North Togo (Savanes and regions). Prefectures had XN rates of 10%. No biochemical data.
Uganda	1991	4139	3.0-4.0 (57.9)			0.40	1656	No national data. A survey in the Kamuli District (1991) found signs of xerophthalmia in 4% of 5074 children between the ages of 0-6 years. Of these, 2.7% of the cases were nightblindness (history), but 1% had X1B indicating a significant public health problem in at least the Northern and possibly Eastern districts of the country. Of those children <5 years of age, approximately 3% showed signs of xerophthalmia. There is a large percentage of children with inadequate dietary intakes resulting in a population where 1/3 of the children are at high risk and 1/5 at medium risk of VAD.

* Refer to Interpreting data in the MDS pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - AFRICAN REGION


Country	Year	Population * (000)	PREVALENCE*			Multiplication factor**	Population at risk** (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
United Republic of Tanzania	1984 1982-85	6036	1.5 (54.3)	45.3		0.60	3622	Large survey (1984) conducted in 8 provinces found a total xerophthalmia rate of 1.53% with VAD a problem of public health significance in Iringa province where the X1B rate was 1.6%. Data from Tabora region (1982-85) indicated that 0.6% of 3177 preschool-age children had X1B and 45.3% had serum levels <0.70 µmol/l.
Zaire		8595						No data available. VAD likely.
Zambia	1988 1985	1790	1.4 (15.0)		16.5	0.60	1074	No national data. A large-scale VAD prevalence survey in Luapula Province (1985) of children <5 years of age found moderate levels of clinical xerophthalmia (XN=1.26%, X1B=0.44%, total xerophthalmia = 1.89%) but revealed indications of past VAD with a XS rate of 0.68%. The biochemical tests found 16.5% of those surveyed had serum retinol levels <0.35 µmol/l. Data available from Ndola Province (1988) show that VAD is a significant public health problem with XN rates as high as 5.0% in rural areas and 13.6% of the total population of 6-12-year-olds having serum retinol levels <0.70 µmol/l.
Zimbabwe	1991	2011	0.6 (4.8)	X ^B		0.40	804	No national data. VAD prevalence data (1991) available on 6 districts indicate a low level of acute clinical xerophthalmia in an unspecified age group. Only a single district surpassed WHO criteria (>1% XN), while two districts had alarmingly high rates of corneal xerophthalmia attaining 0.56%.

* Refer to *Interpreting data in the MDS* pg. 15

Prevalence of Vitamin A Deficiency

Region of the Americas

April 1995



The designations do not necessarily imply VAD severity is uniformly distributed throughout each country

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - REGION OF THE AMERICAS

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Antigua & Barbuda								No data available. No problem likely.
Argentina		3272						Survey in Greater Buenos Aires found that 23.8% of children <1 year of age and 28% of children aged 1-2 year had less than the recommended daily intake of vitamin A.
Bahamas		25						No data available. No VAD likely.
Barbados		20						No data available. No VAD likely.
Belize	1990	34			10.0	0.40	14	Biochemical data available from a national survey (1990) showed that 10% had serum retinol levels <0.70 $\mu\text{mol/l}$, indicating a mild to moderate subclinical VAD problem. Although the surveyed 2-8-year-olds had acceptable serum retinol levels (mean 1.03 $\mu\text{mol/l}$), over 60% had inadequate hepatic stores (RDR > 20%). Evidence points to differences between ethnic groups in risk patterns, i.e. Garifuna and Kekchi Indians (South of country) had highest rates of VAD.
Bolivia	1991	1178			11.3	0.40	471	National data (1991) from the poorest areas of the country indicate that 11.3% of children 1-5 years of age have serum retinol levels <0.70 $\mu\text{mol/l}$, those living in areas of high population density (urban areas) being affected the greatest by severe subclinical VAD. Clinical deficiency was not reported as compared to an earlier national survey (1985) which found a XN rate of 2.3%. Vitamin A capsules have been distributed to those <5 years of age living in high risk areas.
Brazil	1989	16 894		54.7		0.25	4224	Survey in Paraiba (1983) found a moderate to severe rate of X1B (0.5%). Comprehensive biochemical data are available from Coastal Region, Paraiba and the Northeast Region of the country and have consistently demonstrated high prevalences of subclinical VAD. A survey in the Bahia-Semi Arid region (1989) on children aged 0-6 years found that 54.7% had serum retinol levels of <0.70 $\mu\text{mol/l}$. Survey of daily energy intakes found an inadequate consumption of available local foods rich in vitamin A.
Canada	1977	1986						VAD not considered a public health problem.

* Refer to Interpreting data in the MIDS pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - REGION OF THE AMERICAS

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor**	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Chile		1514						Old national data (1960) indicated that 21.6% of children had inadequate vitamin A intake (serum retinol <0.70 µmol/l), but no recent information is available. No evidence of current problem.
Colombia	1977	3875		24.1		0.40	1550	No current data available. Old survey (1960) indicated that 16% of children had serum retinol levels <0.70 µmol/l, while later survey data (1977) documented higher levels still with >20% below this cut-off. No children, however, had serum retinol <0.35 µmol/l.
Costa Rica	1981	420						National data (1981) have demonstrated that VAD is not a severe public health problem, with only 1.8% of preschool children having serum retinol levels <0.70 µmol/l.
Cuba		930						No data available. Subclinical VAD likely.
Dominica								No data available.
Dominican Republic	1991	997	2.8 (11.2)		19.6	0.40	399	No national data. Biochemical survey undertaken in Southwest region bordering Haiti (1991) indicated that 19.6% of preschool children had serum retinol levels <0.70 µmol/l, with 4% showing levels <0.35 µmol/l. A hospital study in Santa Domingo (1991) found a xerophthalmia rate of 2.8%.
Ecuador	1993	1549			16.3 ^b	0.40	620	National survey in five poor provinces (1993) found that 21.9% of those surveyed had serum retinol levels <0.70 µmol/l. National nutrition survey (1986) measured serum retinol in preschool children and noted that 14.1% had serum retinol levels <0.70 µmol/l, with a somewhat higher proportion in rural areas (16.4% <0.70 µmol/l). In dietary surveys, preschool children consumed just over 50% of recommended vitamin A levels, while 81.3% of those children living in rural highlands had dietary patterns which provided <50% of the recommended vitamin A intake. Vitamin A capsules were distributed to the 1-6-year-old population during 1988-1990.
El Salvador	1988	857		36.0		0.40	343	The American Foundation for the Blind conducted a national survey (1973) which found a low prevalence of clinical xerophthalmia (X1B=0.05%), although more recent biochemical data (1988) indicate significant subclinical VAD with 1 out of every 3 preschool children having serum retinol levels <0.70 µmol/l, the highest risk areas being rural.

* Refer to Interpreting data in the MDDS pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - REGION OF THE AMERICAS

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Grenada								No data available. No VAD likely
Guatemala	1986 1988-89	1788		26.0 ^B		0.40	715	Limited clinical data in rural areas (1986) on 0-10-year-olds have indicated a total xerophthalmia rate of 0.5%. However, biochemical data in rural areas collected in 1988 and 1989 have noted as many as 26% of children under 5 years of age with serum retinol levels <0.70 µmol/l. Evaluation of a sugar fortification trial undertaken over 2 years between 1975-77 saw a drop in low retinol levels from 18.7% to 8.3%.
Guyana	1976	94					7	PAHO survey (1976) found no clinical signs of deficiency, though 5.4% of those tested (7.4% in urban areas) had serum retinol levels <0.35 µmol/l.
Haiti	1975	1083	8.0 (52.0)	X		0.60	650	National survey (1975) of an unspecified age group observed a total xerophthalmia rate of 8%, and estimated that as many as 2.5% of the children had corneal involvement. Haiti has a national VAD control program based primarily on capsule distribution, although HKI has also emphasized the production and consumption of vitamin A-rich foods.
Honduras	1987	960			20.0	0.75	720	National nutritional survey (1987) found that 20% of the children had retinol levels <0.70 µmol/l. It also indicated that almost 75% of all families consumed <50% of the recommended vitamin A. VITAL has supported a rapid appraisal of VAD status, but data are not yet available.
Jamaica		266						No data available. No VAD likely.
Mexico	1990	11 912		32.0		0.40	4765	No national data. Data from Hermosilla (1990) indicated that 32% of children 2-7 years of age had serum retinol levels <0.70 µmol/l. Data from Yucatan (1984) indicated clinical deficiency among displaced persons.
Nicaragua	1993	769		31.3		0.40	308	USAID preliminary biochemical results (1993) found that 7.9% of the population studied was deficient (<0.35 µmol/l) and that 31.3% had marginal serum levels (<.70 µmol/l). Old national data (1966) indicate approximately 20% of children aged 0-4 years had serum retinol levels <0.70 µmol/l. A nutritional study (1982) in the northern and central areas found that children aged 7-12 years consumed <10% of the recommended amount of fruits and vegetables. No recent biochemical studies done.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - REGION OF THE AMERICAS

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Panama	1992	309						National data (1992) indicated mild levels of VAD with 6.0% of children under 15 years of age having serum retinol levels <0.70 µmol/l. VAD is not a problem at the public health level, but it appears that preschoolers are at risk of inadequate vitamin A intakes, especially indigenous people who tend to have lower retinol levels.
Paraguay	1965	716						Old national data (1965) indicate that approximately 12% of children under 14 years of age had serum retinol levels <0.70 µmol/l. No recent data available.
Peru	1992	3010		22.0*		0.40	1204	No clinical data available. Biochemical data from Piura and Puno provinces (1992) indicate a high proportion of inadequate vitamin A intake (retinol levels <0.70 µmol/l) in 32.8% of children 0-6 years of age and 14.1% of preschool children, respectively. Other information on VAD from older dietary surveys (1970s) note that almost all families surveyed in Puno province consumed <50% of the recommended level of vitamin A. Other provinces of Piura, Junin and Lima-Callao also had marginal vitamin A intakes.
Puerto Rico		315						No data available.
St. Kitts & Nevis								No data available. No VAD likely.
St. Lucia								No data available. No VAD likely.
St. Vincent & the Grenadines								No data available. No VAD likely.
Suriname		55					7	No data available.
Trinidad & Tobago		144						No data available. No VAD likely.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - REGION OF THE AMERICAS


Country	Year	Population * (000)	PREVALENCE*			Multiplication factor**	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
United States of America	1970s- 1980s	20 344						Data from Health and Nutrition Examination surveys (1970s-1980s) and studies of Mexican Americans, Puerto Ricans and African Americans (1970s-1980s) indicate mean serum retinol levels between 1.04 and 1.31 $\mu\text{mol/l}$. Data from Mexican Americans in the southwest (1982-83) found that only 4.6% had serum levels $<0.70 \mu\text{mol/l}$ and that non-Hispanic blacks and whites had mean serum levels of 1.04 $\mu\text{mol/l}$ and 1.15 $\mu\text{mol/l}$, respectively. Only 10% of those surveyed (1982-83) had serum retinol levels $<0.70 \mu\text{mol/l}$.
Uruguay	1962	262						Old data from a nutritional survey (1962) showed no evidence of clinical or biochemical hypovitaminosis A.
Venezuela	1960	2566						Old national data (1960) indicated that between 5-8% of children <14 years of age had serum retinol levels $<0.70 \mu\text{mol/l}$, but no recent data on vitamin A status are available.

* Refer to *Interpreting data in the MDS* pg. 15

Prevalence of Vitamin A Deficiency

South-East Asian Region

April 1995



SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - SOUTH-EAST ASIA REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Bangladesh	1983	19 633	4.6 (678.8)			0.75	14 725	A large-scale national blindness survey (1982-83) provides the most up-to-date estimate of VAD in the country. Rural areas had the following rates of clinical xerophthalmia: XN=3.6%, XIB=0.9%, corneal involvement =0.1%. Urban slum areas: XN=2.8%, XIB=1.6%, corneal involvement =0.2%. The total xerophthalmia rate was 4.6%. The sample design was proportional to the population distribution, found primarily in rural areas, and thus was similar to the overall rates which were: XN=3.5%, XIB=1.0%, corneal involvement=0.1%. This translates to 11 cases per 10 000 or 11 times greater than WHO considers to be a severe public health problem. Biochemical data from Matlab (1980) found 20% of preschool children to have serum retinol levels <0.35 µmol/l. VAD in Bangladesh is linked to the heavy dependence on a diet of rice. Surveys have showed low levels of vitamin A intake on a national basis and efforts are under way to improve dietary diversity through increased production and consumption of vitamin A-rich foods.
Bhutan	1989	277	0.7 (1.4)	X		0.75	208	A large-scale national nutrition survey (1989) was undertaken during which signs of clinical xerophthalmia were noted. A total xerophthalmia rate of 0.7% among children aged 0-5 years confirmed an earlier observation (1976) of moderate to mild clinical xerophthalmia. However, a national biochemical study (1985) noted that 14% of preschool children and 13% of pregnant women had serum retinol levels <0.35 µmol/l indicating a significant subclinical VAD problem. Dietary information from 1981-82 shows that 88% of households consume less than the RDA.
Democratic People's Republic of Korea		2687						No data available. No VAD likely.

* Refer to *Interpreting data in the AIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - SOUTH-EAST ASIA REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
India	1988-90	115 894	0.7 (608.4)			0.75	86 920	Over the past 10-15 years VAD rates have been monitored by the National Nutrition Monitoring Bureau (NNMB) which have noted that the prevalence of X1B has declined between 1976 and 1990 from 2.0 to 0.7%. One survey (1988-90) found a X1B rate of 0.7%. A baseline survey (1989) for a vitamin A supplementation trial in Tamil Nadu noted high xerophthalmia rates, including XN=3.7%, X1B=7.2% and total xerophthalmia rate = 10.95%. Biochemical data from the same survey found that 37.5% had retinol levels <0.70 µmol/l. India has a comprehensive strategy to combat VAD, primarily through vitamin A supplementation. Home gardening and education programs have been initiated in three states.
Indonesia	1991	23 532		50-65		0.60	14 119	Over the past fifteen years, most of the pioneering work in vitamin A control programme development and research has taken place in Indonesia, including evaluation of the mortality effects of vitamin A supplementation and trials with MSG fortification. A large-scale national prevalence survey (1977-78) of children aged 0-5 years found an overall X1B prevalence of 1.0%, corneal xerophthalmia of 0.06% and XS of 0.13% indicating a severe problem in the past. Between 1983 and 1990, many of the provinces earlier identified as having the highest xerophthalmia rates were re-surveyed, and considerable reduction were noted. Later studies in several provinces (1991) were showing total xerophthalmia rates of <1.0%. However, biochemical studies showed that >50% of those surveyed had serum retinol levels <0.70 µmol/l. Vitamin A supplementation activities have been extended to include home gardening and education to promote the consumption of vitamin A-rich foods.
Maldives	1977	42						A country report on Felidu Island (1977) of children <6 years of age found no cases of xerophthalmia, though results seem to be dependent on the availability of fish, liver and greens.
Mongolia		371						No available data.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - SOUTH-EAST ASIA REGION


Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Myanmar	1987-89	6509	2.0 (78.1)	32.4 ^B		0.60	3905	No national data. Large-scale study undertaken in central Myanmar (1978-1990) noted some high levels of X1B ranging from 0.7% in Natogyi to 3.3% in Saku (though considerably smaller sample size in latter region). A survey conducted in dry upper zone (1987-89) of 4 regions observed X1B of 2.0% among children aged 2-5 years, but somewhat higher rates in school-age children, including >9% of 10-14-year-olds in Monywa. Biochemical data from the same study noted a similar range in serum retinol values with as much as 76.1% of children aged 2-6 years living in Kyaukpadaung having serum retinol levels <0.70 µmol/l and over 16% with values <0.35 µmol/l.
Nepal	1981 1993	3442	3.0 ^A (77.5)			0.75	2582	Nepal had a large-scale blindness xerophthalmia survey (1981) in which the overall prevalence of X1B among children aged 0-6 years was 0.64%. However, there was considerable variation in the distribution within the country, with the highest rates observed in the Terai region along the border with India. A large intervention study in the Central Terai noted X1B prevalences above 2% (districts of Bara and Parsa at baseline in 1989). A survey undertaken in the Jumla district (1990) noted a 13.2% prevalence of active xerophthalmia in preschool children. Surveys in the far and midwest (1993) found X1B rates around 3.0%.
Sri Lanka	1987	1793	0.3-0.7 (3.7)			0.40	717	A national prevalence survey (1987) an average X1B prevalence of 0.3%. Of the districts surveyed, Kegalle and Kurunegala had an average X1B prevalence of 0.70%. Subsequently, the Ministry of Planning and Implementation published VAD prevalence data for all districts which indicate the problem may be even more widespread.
Thailand	1990 1992	5576			20.0	0.25	1394	No national data. Survey in the south (1992) found a XS rate of 0.90% and corneal involvement of 0.67%. Problem seemed to be isolated to south due to use of unfortified condensed milk during infant feeding. Data available from Sakon Nakhon province (1990) in the northeast part of the country reported a 0.0-1.3% prevalence of XN among children aged 2-6 years in rural areas, depending on season. An important seasonal difference has been noted in the vitamin A levels of preschool children in the north and northeast, with serum retinol means being significantly higher in the months of September and October (1.3 µmol/l) when compared to those found in February and March (1.02 µmol/l). MOH reports no clinical cases since early 1994.

* Refer to *Interpreting data in the MIDS* pg. 15

Prevalence of Vitamin A Deficiency

European Region

April 1995



The designations do not necessarily imply VAD severity is uniformly distributed throughout each country

WHO 95219

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EUROPEAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Albania		358						Rapid assessment of eight districts (1991) found little evidence of VAD.
Armenia		351						No data available.
Austria		454						No data available.
Azerbaijan		787						No data available.
Belarus		596						No data available.
Belgium		603						No data available.
Bosnia and Herzegovina		220						No data available.
Bulgaria		449						No data available.
Croatia		249						No data available.
Czech Republic		1094						No data available.
Denmark		323						No data available.
Estonia		83						No data available.
Finland		319						No data available.
France	1985-86	3826						Study in Tours (1985-86) of 1-16-year-olds recorded a serum retinol mean of 1.47 $\mu\text{mol/l}$. Nutritional Study on 0-3-year-olds throughout twenty day-care centres in Paris found that more than 70% of the vitamin A RDA was received while at day care.
Georgia		417						No data available.
Germany		4571						No data available.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EUROPEAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Greece		526					No data available.	
Hungary		636					No data available.	
Iceland		23					No data available.	
Ireland		248					No data available.	
Israel	1970s	582				?	No national data. A small survey (1970s) of low-income Arab children living in East Jerusalem noted that 83% of girls and 86.9% of boys had serum retinol levels <0.70 µmol/l with mean serum retinol values of approximately 0.60 µmol/l.	
Italy		2883					No data available.	
Kazakhstan							No data available.	
Kyrgyzstan							No data available.	
Latvia		181					No data available.	
Lithuania		273					No data available.	
Luxembourg		23					No data available.	
Malta		27					No data available.	
Monaco							No data available.	
Netherlands		1048					No data available.	
Norway		317					No data available.	
Poland		2708					No data available.	

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EUROPEAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Portugal		564					No data available.	
Republic of Moldova		339					No data available.	
Romania	1970s	1775					Only data available are from a survey (1970s) of 117 orphans which resulted in a mean serum level of 0.74 $\mu\text{mol/l}$ (± 0.33).	
Russian Federation		22 845					No data available.	
San Marino							No data available.	
Slovakia		373					No data available.	
Slovenia		102					No data available.	
Spain		2101					No data available.	
Sweden		606					No data available.	
Switzerland		433					No data available.	
Tajikistan		986					No data available.	
The former Yugoslav Rep. of Macedonia		158					No data available.	
Turkey	1985-86 1994	7779					No national data available. Small study done in Marmara in central and east Anatolia (1986) of 150 children aged 7-17 years. Biochemical measures were taken and an overall mean serum retinol level of 1.18 $\mu\text{mol/l}$ (± 0.46) was observed, with consistent levels across age groups and no differences by sex. Another survey (1994) of 56 children receiving measles immunizations found that 9.5% of the healthy, 42.9% of sick controls, and 90.5% of the measles group had serum retinol levels $<0.70 \mu\text{mol/l}$.	

* Refer to Interpreting data in the MIDS pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EUROPEAN REGION


Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Turkmenistan		577					No data available.	
Ukraine		2884					No data available.	
United Kingdom of Great Britain and Northern Ireland		3969					No data available.	
Uzbekistan	1993	3222		X			Data from Muynak District (1993) found that 48.9% of children aged 5-6 years had serum retinol levels $<0.70 \mu\text{mol/l}$.	
Yugoslavia		1645					No data available.	

* Refer to *Interpreting data in the MDS* pg. 15

Prevalence of Vitamin A Deficiency

Eastern Mediterranean Region

April 1995



SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EASTERN MEDITERRANEAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Afghanistan	1989	4250		X			?	Study in Kabul (1989) indicated a 2.4% X1A rate among children between the ages of 5-14 years. No biochemical data are available.
Bahrain		69						No data available. VAD not considered a public health problem.
Cyprus		60						No data available. No VAD likely.
Djibouti	1988	95			16.0 ^E	0.40	38	Comprehensive survey (1988) of children aged 0-5 years available including biochemical and clinical parameters for rural and urban areas. The prevalence of X1B was found to be 1.04% in rural areas, while in urban areas it was 0.43%. However, biochemical data on children aged 4-10 years gave a different impression, with 12.3% of rural children having serum retinol levels <0.70 µmol/l, but over 20% of urban children falling below the 0.70 µmol/l cut-off. These data were further confirmed by CIC (with transfer) where 13.6% of the urban children had morphology indicative of VAD, while only 8.3% of the rural children were classified as vitamin A deficient.
Egypt	1978	7798					?	Very few studies exist. Survey by MOH (1978) found VAD among 0.5% of children sampled. Though small, unrepresentative assessments do not suggest a VAD problem, the UNICEF regional office (Amman) reports that it is.
Iran (Islamic Republic of)	1980s	11 589			X		?	No national data. In a UNICEF report <i>Situation of Women and Children in Iran</i> , several survey data were summarized including biochemical information from 4 provinces (1980s) where 9% of preschool children had serum retinol levels <0.70 µmol/l. In studies in Kohguituyeh and Boyerahmad, no boys aged 7-17 years had serum retinol levels <0.35 µmol/l. In a goitre endemic area of Teheran, no girls surveyed had serum retinol <0.70 µmol/l.
Iraq	1994	3574	1.6 (14.3)	X ^E		0.25	293	No national data available. UNICEF preliminary data from health centres in Baghdad, Basrah and Mosul (1994) found a XN rate of 1.6%. In a rapid situational assessment in Hartha (1992) following the Gulf War, a 2.9% XN rate and 1.4% X1B rate were observed in preschool children (95% CI 0.2-5.7) while an examination of 16 malnourished children in Basrah yielded one case of Bitot's spots. Furthermore, there is a high incidence of diarrhoea (52.5-59.8%) and respiratory disease (2.7-5.5%).

* Refer to *Interpreting data in the MIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EASTERN MEDITERRANEAN REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor**	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Jordan		832					?	Old survey (1960) noted a 1.27% XN rate. Nutritional survey (1975) of children aged 0-5 years showed a clinical prevalence of 0.05%. No current data, though the UNICEF regional office reports that VAD is of public health significance.
Kuwait		223						No information available. UNICEF regional office believes that VAD is not a serious problem in the country.
Lebanon		376					?	Data not yet available from a survey conducted in late 1994. An old nutrition survey (1962) showed that 3% of children aged 5-9 years had serum retinol levels <0.35 µmol/l. No signs of Bitot's spots were seen in 98 children aged 0-4 years. UNICEF regional office considers VAD to be likely.
Libyan Arab Jamahiriya		959					?	No data available. UNICEF regional office considers VAD to be likely.
Morocco	1971	3955					?	Old national data (1971) indicates a 1% total xerophthalmia rate for children under four years of age. No recent data available.
Oman	1994	328			20.8	0.60	197	National survey (1994) found that 20.8% of those surveyed had serum retinol levels <0.70 µmol/l, indicating a moderate to severe subclinical problem. Children 18 months of age were affected the most with 22.8% having serum retinol levels <0.70 µmol/l. Data from 12 MCH Centres (1981) noted X1B prevalences to be 1.5% with higher rates in boys than in girls. However, more recent data (1991) on children aged 0-2 years found no cases of clinical xerophthalmia, indicating that clinical VAD may be under control. Supplementation programs have been proposed for 1995.
Pakistan	1990 1988-89	22 550	0.0	29-70*		0.40	9020	Clinical xerophthalmia observed in several small studies in Lahore and West Pakistan in 1960s. A national micronutrient survey (1976-77) reported a X1B rate of 1.5% in preschool children. Biochemical data from the same survey noted that 13.0% of school-age children had serum retinol levels <0.70 µmol/l, although it was noteworthy that only 2.1% of lactating mothers had marginal serum vitamin A. A survey in Orangi (1987) found a X1B rate of 3.8%. Smaller surveys (1988-89) in several urban slums of Karachi (Azambasti, Chenesar-Goth and Esanagri Gilgit) found no signs of xerophthalmia though the biochemical data indicate that VAD is a public health problem with 29-70% of preschool children having serum retinol levels <0.70 µmol/l.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - EASTERN MEDITERRANEAN REGION


Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Qatar		51						No data available. UNICEF regional office reports that VAD is unlikely.
Saudi Arabia	N.S.	2822						No national data. Results of a biochemical study in Riyadh indicate that 10% of the population surveyed had serum retinol levels <0.70 µmol/l and that 1.1% had serum values <0.35 µmol/l, indicating that VAD is not a severe problem.
Somalia	1993	2003	1.5 ^a (7.5)		X ^b	0.25	500	WHO's National Morbidity Survey (1982) showed that hypovitaminosis A was not a public threat. Reports of VAD occurred after the Civil War in central, southern, northwest and northeast zones. Nutritional surveys in central Somalia report 1-2% of X1B.
Sudan	1986-87 1989	4989	1.6 ^a (59.9)			0.75	3742	No national data. However, clinical surveys conducted in many of the provinces have indicated that VAD is a significant public health problem. Studies (1986-87) found XN prevalence rates ranged from approximately 0.5% in Gezira, Kassala, Nile and Northern areas to above 2% in Khartoum, Darfur and Kordofan. Studies done in refugee camps have observed alarmingly high levels of clinical xerophthalmia, but these were all based on very small sample sizes. In the Red Sea Province (1989), xerophthalmia is around 1.6%. Food consumption survey in Upper Nile (1978) found that a normal diet contained only 6-10% of RDA. No biochemical data available.
Syrian Arab Republic		2746						No data available. UNICEF regional office considers VAD to be likely.
Tunisia		1089						No recent data available. A nutritional survey (1975) concluded that hypovitaminosis A was not a problem, though UNICEF regional office considers VAD to be likely.
United Arab Emirates		177						No data available. UNICEF regional office considers VAD to be likely.
Yemen	1992	2705	2.1 (42.6)	62.4		0.75	2029	Preliminary data from 21 villages in the Tihama region (1992) observed an overall X1B rate of 1.7% and a total of 2.14% in children aged 1-6 years. It is noteworthy that the prevalence of X1B was directly related to age, with the highest X1B rates seen in children 5-6 years of age (2.94%). It was found that 62.4% of children aged 1-5 years showed serum retinol levels <0.70 µmol/l. Survey of health centres (1981) found no keratomalacia, acute corneal xerosis or ulceration.

* Refer to *Interpreting data in the MDS* pg. 15

Prevalence of Vitamin A Deficiency

Western Pacific Region

April 1995



The designations do not necessarily imply VAD severity is uniformly distributed throughout each country

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - WESTERN PACIFIC REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Australia		1352						No data available. No VAD likely.
Brunei Darussalam		32						No data available. No VAD likely.
Cambodia	1993	1505	6.9 (77.9)			0.75	1129	No national data. Survey of 5 provinces (1993) found a XN rate of 6.3% and a X1B rate of 0.6%
China	1982	120 386			18.5*	0.25	30 096	National data from urban Beijing (1982) found 17% of 0-2 year olds had serum levels <0.70 µmol/l. Study of rural children notes that as many as 20% of children between 6 months and 2 years of age have serum retinol levels <0.70 µmol/l. Dietary surveys in rural areas show that children <3 years of age consume only 30-40% of the RDA of vitamin A. Xerophthalmia and blindness caused by VAD is rare. Largest XN rate was found in Talep (11.0%) and Kompong Thom (9.8%).
Cook Islands	1989-92		6.0				?	Surveys done in high-risk malnutrition areas to determine severity of VAD. Results found no cases of xerophthalmia.
Fiji		84						No data available. No VAD likely.
Japan		6912						National nutritional survey (1988) of 20 000 people showed an intake of 2596 IU per capita per day.
Kiribati	1994		1.1 (?)			0.75	?	Survey of three villages on Kiritimati Island (June 1994) found a xerophthalmia rate of 1.12% (X1A, X1B) among 0-6-year-olds. However, national data from the Gilbert islands (1989) indicate that VAD may be quite severe, with 14.7% of preschool children aged 6 months to 5 years with xerophthalmia, 10.9% from X1B. Short-term vitamin A supplementation programme is almost completed.
Lao P.D.R.	1995	887		X		0.40	335	National survey (1995) found that among children ages 24-71 months, 0.7% reported nightblindness. One child reported Bitot's spots and another had an active corneal ulcer. Low intake of dark green leafy vegetables. No other data are available.

* Refer to *Interpreting data in the MIDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - WESTERN PACIFIC REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Malaysia	1984	2656			12.0	0.25	664	No national data. Data from the coastal and river area (1978) indicate cases of VAD, especially in Sarawak River Delta where a xerophthalmia rate of 38.2% was found. Biochemical data (1984) from a rural area in peninsular Malaysia found that 12% had serum retinol levels <0.70 µmol/l. However, severe problem seems to be confined to certain areas such as rural areas and does not seem to pose a public health problem.
Marshall Islands	1991		4.0			0.75		UNICEF survey (1991) found a xerophthalmia rate of 4.0%.
Micronesia (Federated States of)	1988-89		20 (?)	64.0		0.25	?	National data available from Chuuk (1988-89) found an alarmingly high prevalence of xerophthalmia with XN rates of 14.0% and X1B rates of 6.0% in children 3-7 years of age. Subsequently, serum retinol levels of these children were found to be very low with 15% <0.35 µmol/l and 64% <0.70 µmol/l.
Nauru								No data available.
New Zealand		300						No data available.
Niue								No data available.
Papua New Guinea	1993 1990s	644	0.6 (1.0)	91.0		0.25	161	Study done in hospitals throughout 9 provinces (1993) found that 0.59% suffered clinically from VAD. VAD suspected to exist from available data in East Sepik where 91% had serum retinol levels <0.70 µmol/l.
Philippines	1993	9421	0.4 (15.3)			0.40	3768	National nutritional survey (1993) found a XN prevalence of 0.7% among children 6 months to 6 years of age. A steady decrease has been documented through the past decade by surveys undertaken in 1982 and 1987 which found national estimates of XN decreasing from 3.5% to 0.9%. Biochemical results from a national vitamin A deficiency prevalence survey (1987) found that 2.6% of the preschool children had serum retinol levels <0.35 µmol/l. Studies in Samar and Zamboanga (1991) indicate that VAD is still highly prevalent in some areas, with X1B rates of over 2% observed in preschool children aged 0-5 years.
Republic of Korea		3548						No data available. No VAD likely.
Samoa								No data available. No VAD likely.

* Refer to *Interpreting data in the MDS* pg. 15

SUMMARY OF VITAMIN A DEFICIENCY PREVALENCE DATA - WESTERN PACIFIC REGION

Country	Year	Population * (000)	PREVALENCE*			Multiplication factor*	Population at risk* (000)	Notes
			Clinical	Severe sub- clinical	Moderate sub- clinical			
Singapore		219					No data available. No VAD likely.	
Solomon Islands	1991	64	1.5 (0.7)			0.75	48	Data from 7 islands (1991) noted an average X1B prevalence of 1.42% among preschool children and a total xerophthalmia rate of 1.52% in children aged 0-5 years.
Tokelau								No data available.
Tonga								No data available.
Tuvalu	1991		0.0				7	National survey found no cases of xerophthalmia.
Vanuatu	1991	27	0.05			0.75		National survey found corneal xerosis rate of 0.05%.
Viet Nam	1988	9630	0.65 (34.7)			0.60	5772	National data (1994) reveals that xerophthalmia is no longer a problem in the country. Among children <5 years of age, a prevalence of 0.14% was found. Earlier national data (1988) found an overall xerophthalmia prevalence to be around 0.65% of which a majority of the clinical symptoms and signs noted were XN (0.35%). However, the rate of corneal xerophthalmia was 0.13%, well above the criteria set by WHO to classify countries as having a significant VAD public health problem. FAO has implemented a number of home gardening studies which are being evaluated.

* Refer to *Interpreting data in the MDS* pg. 15

MDIS

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY

DISAGGREGATED DATA TABLES

GUIDELINES FOR USING DISAGGREGATED TABLES

Information on VAD prevalence in this section is presented in two sets of tables:

- Ocular signs and symptoms
- Serum retinol data

For each set, data are presented for countries in alphabetical order according to the six WHO regions. For each country there is a brief description of the location and population under study in the column marked *Geographic area*, together with any other defining characteristics of the region or area. The *Survey year* refers to the year in which the study was conducted or ended if it spans more than one year. *N.S.* (not specified) is recorded if the survey year is unknown. The *Age group* is presented in years. *Sex* is divided into three categories each denoted by the following letters: B=both males and females; M=males only; F=females only. If the sex was not specified it is assumed that both sexes were assessed, and a 'B' is recorded on the table. *Sample size* refers to the number of children in a particular strata. If the sample size was not given, *N.S.* is inserted. Where no data are given in one or more of these columns for a particular survey, unless otherwise noted it is assumed that it is the same as the preceding survey.

For clinical data, point prevalence estimates are provided for individual signs and symptoms of clinical xerophthalmia and the total of corneal xerophthalmia (X2+X3A+X3B). Where a total xerophthalmia rate is available, the next column indicates the particular classification scheme used, i.e. the specific stages of xerophthalmia which are included in the estimate. It is very important to recall when comparing data across studies that disparate classifications were employed in many of the rates presented for total xerophthalmia

For serum retinol data, in addition to those characteristics outlined above, a column has

been included to specify the laboratory method employed in the biochemical analysis and determination of serum retinol levels. The information for serum retinol is presented both as prevalence, i.e. percent of population falling below 0.70 and 0.35 $\mu\text{mol/l}$, as well as for the entire distribution, by designating the mean value and its standard deviation.

Reference numbers are coded according to WHO region where the first number corresponds to the following: 1=AFR; 2=AMR; 3=EMR; 4=EUR; 5=SEAR; 6=WPR; 7=Multiple country data. Specific references are listed in a separate section of the document.

**PREVALENCE OF OCULAR SIGNS AND
SYMPTOMS**

- DISAGGREGATED TABLES -

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.
						XN	X1A	X1B	X2	X3A	X3B	X5	Corneal	Total *		
AFRICAN REGION																
ANGOLA	Andulo, Bic Province	1973	0-4.99	B	N.S.			3.00						Z	7018	
BENIN	Atacora region	1975	N.S.	B	1965			0.55	2.60		0.20			3.35	D	7001
	North Atacora	1989	0-4.99		1551	3.40		4.90							1041	
BURKINA FASO	4 northern regions	1984	0-5		2786	2.10		12.00				0.00	0.30	14.40	B	7001
	Bam		16-45		1722	2.50		1.90				0.10	0.60	5.00	Z	1035
			2-10		1263	0.36										
			2-5		429	1.16										
			6-10		824	0.35										
	Gourma		2-10		1609	0.87										
			2-5		885	0.79										
			6-10		724	0.96										
	Passore		2-10		1260	1.58										
			2-5		664	1.80										
			6-10		596	1.34										
	Soun		2-10		1055	2.55										
			2-5		583	2.57										
			6-10		472	2.54										
	Tapoa		2-10		948	3.58										
			2-5		483	1.65										
			6-10		465	5.59										
	Yatenga		2-10		2628	0.60										
			2-5		1520	0.32										
			6-10		1108	0.99										
	Districts of Yatenga, Passore, Sourou	2/86	0-5.99		1103	2.82		0.27	0.00			0.09	0.18	3.27	B	7015
				F	N.S.	3.37		0.19	0.00			0.00	0.00	3.56		
				M	N.S.	2.33		0.35	0.00			0.18	0.35	3.03		
			6-10	B	932	2.29		2.17	0.00			0.360	0.00	4.46		
				F	N.S.	2.10		1.47	0.00			0.00	0.00	3.57		
				M	N.S.	2.63		2.85	0.00			0.63	0.00	5.48		
	Passore - rural		0-5	B	107	0.00		0.00	0.00			0.00	0.00	0.00		
			6-10		94	1.55		2.13	0.00			0.00	0.00	3.68		
	- urban		0-5		95	4.41		0.00	0.00			0.00	0.00	4.41		
			6-10		104	2.88		1.92	0.00			0.00	0.00	4.80		
	Sourou - rural		0-5		226	2.98		0.00	0.00			0.00	0.00	2.98		
			6-10		177	5.10		0.56	0.00			0.00	0.00	5.66		
	Yatenga - rural		0-5		675	2.22		0.52	0.00			0.17	0.34	3.08		
			6-10		557	1.53		2.69	0.00			0.66	0.00	4.22		
	3 northern regions	1987	N.S.		N.S.	5.12								13.2	Z	7001
	Six provinces	1989	2-10			1.42		0.26	0.03			0.03	0.05	1.73	B	
CAMEROON	Extreme Northern Province	4-5/92	0-5	B	5352	0.19		0.47			0.12			0.71	Z	7023
	Flooded plains				1644	0.00		0.43						0.55		
	Mountains				1986	0.35		0.65						1.00		
	Plains				1163	0.26		0.26						0.52		
	Periurban				559			0.36						0.54		
CAPE VERDE	San Vicente and Fogo	1982-83	4-70	F	616		3.40	2.90	0.00	0.00					Z	7016
				M	535		5.00	3.60	0.00	0.00						
CHAD	Rural areas	1984-86	0-4	B	1626			0.37			0.06	0.49			Z	1002
			5-9		1379			0.94			0.07	0.65				
	- drought camps		0-9		180			1.67			1.11	0.00				
	- equipped camps				559			0.00			0.00	0.72				
	- sedentary				2263			0.71			0.00	0.57				
	Subnational	1984	0-5		N.S.		2.20	0.50	0.00			0.40			Z	7001
	Chari-Baguirne, Bath, Ouaddi	3/86			1044	2.18		0.48	0.00			0.38	0.00	2.66	B	7015
				F	901	2.44		0.37	0.00			0.19	0.00	2.81		
				M	809	1.89		0.59	0.00			0.59	0.00	2.48		
			6-10	B	665	1.65		1.65	0.00			0.90	0.15	4.35		
				F	N.S.	0.82		1.37	0.00			0.82		2.19		
				M	N.S.	2.68		2.00	0.00			1.00	0.33	5.01		
	Batha - rural		0-5	B	481	1.37		0.21	0.00			0.42	0.00	1.58		
			6-10		318	1.89		1.57	0.00			1.57	0.31	3.77		
	Chari-Baguirne - rural		0-5		377	3.48		1.06	0.00			0.53	0.00	4.54		
			6-10		214	1.87		0.47	0.00			0.47	0.00	2.34		
	- urban		0-5		71	2.08		0.00	0.00			0.00	0.00	2.08		
			6-10		32	0.00		0.00	0.00			0.00	0.00	0.00		
	Ouaddi - rural		0-5		115	1.27		0.00	0.00			0.87	0.00	1.27		

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms								Total *	Ref. No.		
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal				
ETHIOPIA	Rural areas total		6-10		101	0.99		5.00	0.00			0.00	0.00	5.99			
			0-5		973	2.19		0.51	0.00			0.00	2.70				
			6-10		631	1.56		1.56	0.00			0.99	0.14	3.26			
	National	1980-81	5-6.99	B	6636		4.80	1.00						5.90	C	1052	
	- Cropping zone				3827		5.90	1.10						7.10			
	- Cash crop zone				1114		1.90	0.40						2.30			
	- Ensete zone				480		0.80	0.00						0.80			
	- Pastoral zone				1215		5.30	1.60						6.90			
	Shoa - 7 Provinces:	1981	6-20	B	14 740			0.91				1.40				1065	
				F	5961			0.60				1.00					
				M	8779			1.10				1.60					
			6-10	F	2342			0.50				0.50				7016	
				M	3954			1.00				1.80					
			11-14	F	3182			0.70				1.30					
				M	4068			1.20				1.80					
			15-18	F	426			0.90				1.60					
				M	705			1.60				2.70					
		Chebona-Gurage		6-20	B	5162			0.30								
				F	1457			0.10									
				M	3705			0.40									
		Kembatana Hadya			B	1836			0.30								
				F	773			0.30									
				M	1063			0.40									
		Menzena-Gishe			B	405			0.20								
				F	167			0.60									
				M	238			0.00									
		Merhabete			B	1333			1.80								
				F	629			1.40									
				M	704			2.10									
		Selate			B	867			2.30								
				F	374			1.10									
				M	493			3.20									
		Teguletua-Bulga			B	1374			0.60								
				F	676			0.40									
				M	698			0.70									
		Yifatna Timuga			B	3763			1.60								
				F	1878			2.30									
				M	1885			0.80									
		Sudan: Wad Kowli - Ethiopian refugees	1985	<10	B	929		0.11	2.48	1.61			2.48		6.70	Z	1006
		Sudan: Wad Sherife - Ethiopian refugees				262		0.00	1.91	1.91			2.29		6.10		
		Camp Adi Gafuf - supplementary feeding		0-14.99		427		0.50	1.60				0.50	0.70	2.80	C	1029
		Camp Aji Bar		0-12.99		60		3.30	3.30				1.70	0.00	6.60		
		- dry feeding															
		- intensive feeding				451		0.90	2.90				1.80	0.90	4.70		
		- super intensive feeding				110		0.00	0.90				0.90	0.00	0.90		
		- total				621		1.00	2.60				1.60	0.60	4.20		
		Camp Ansokia (Wello)		0-14.99		37		0.00	0.00				0.00	0.00	0.00		
		- super intensive feeding															
		- intensive feeding				228		4.80	1.80				0.00	0.40	7.00		
		- total				265		4.20	1.50				0.00	0.370	6.07		
	Camp Bete		0-9.99		167		1.20	3.00				1.20	0.60	4.80			
	- general feeding				237		0.40	1.70				0.40	0.40	2.50			
	- monthly feeding				321		0.60	1.90				0.30	0.00	2.50			
	Camp Quiba - supplemental feeding		0-14.99		319		2.80	0.90				0.90	0.60	4.30			
	Arsi and Bale		5-6.99		2647		16.36	3.89				0.42	0.60			1004	
	Arsi				1241		20.47	6.29				0.64	0.73				
	Bale				1406		12.73	1.78				0.21	0.50				
	Debara School: Arsi	1987	7-16		175		29.00									7016	
				F	43		18.30										
				M	132		32.50										
	Illo: Arsi		2-15	B	475		43.60										
				F	212		39.10										
				M	263		47.10										
	Bahir Dar School: Gojjun		4-6	B	538		1.85										
				F	272		1.83										
				M	266		1.87										
	Juna School: Keffun		5-6	B	706		1.27										
				F	363		1.60										
				M	343		0.87										

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.			
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *					
GHANA	Melkaye village: Hararge -with signs of xerophthalmia -without signs of xerophthalmia	1990s	0-12.5	B	240	28.75		6.67	0.83			5.83	6.25		1073				
				F		12.10													
				M		12.90													
				F		25.00													
GHANA	Kassena-Nankana VITAL Survival Study - Kassena-Nankana VITAL Morbidity Survey Kassena-Nankana	1989-91	0-4 .5-7.5	B	16 568	1.04		0.02	0.01			0.05		Z	7016				
					21 906	0.70								1064					
		1991			1177	1.50						0.20			7016				
GUINEA	North Guinea - pregnant females - lactating females	N.S.	pregnant	F	N.S.			5.20						Z	1040				
KENYA	National Data Baringo - Highland Tribe - Lowland Tugen - Njemps - Pokot Kajiado - Maasai Tribe Kakamega - Abaluhya Kisii - Kisii Tribe Kwale - Giriama Tribe Meru - Meru Tribe Nyanza - Luo Tribe Nyeri - Kikuyu	1976-81	N.S.	B	13 781	0.05	0.21	0.00	0.00	0.00	0.00	0.09	0.00	0.26	C	1039			
					702	0.00	1.13	0.00	0.00	0.00	0.00	0.00	1.13						
					472	0.21	0.36	0.00	0.00	0.00	0.21	0.00	0.57						
					590	0.00	0.00	0.00	0.00	0.00	1.01	0.00	0.00						
					595	0.33	0.84	0.00	0.00	0.00	0.33	0.00	1.17						
					1920	0.00	0.05	0.00	0.00	0.00	0.16	0.00	0.05						
					1649	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.06						
					1753	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00						
					1339	0.07	0.22	0.00	0.00	0.00	0.00	0.00	0.29						
					1137	0.11	0.61	0.00	0.00	0.00	0.00	0.00	0.72	Z					
					1807	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00						
					1817	0.05	0.05	0.00	0.00	0.00	0.00	0.00	0.10						
LESOTHO	National Micron. Survey	1993	2-6	B	249			0.00					0.00	Z	1055				
MALAWI	Lower Shire - dry season	1983	0-5.99	B	5436	3.39	0.09	0.33					0.05	3.86	A	1021			
					5436	5.40	0.09	0.33				0.06	5.88		7004				
					2818	3.40	0.10	0.21				0.07	3.80	Z	1021				
					2597	3.41	0.07	0.46				0.04	4.05						
						N.S.	0.00	0.00				0.00	0.00	A					
							0.00	0.10				0.10	0.20						
							0.00	0.21				0.00	3.71						
							4.56	0.20	0.41			0.10	5.27						
							7.15	0.20	0.74			0.06	7.41						
							N.S.						3.90	Z	1018				
													0.55		1033				
						Mbalachanda	7/88	0-4		2542	0.12	0.43			0.00				
						Salima and Dedza	9/88	2-7		650	1.40	0.50	0.20			0.00	0.00		7016
	Mkhota	3/89	0-8		3791	0.80	0.60	0.20	0.03		0.08	0.03	1.66	1033					
MALI	Segou region North-following severe drought Sikasso - rural - urban Displaced population Nomadic population Sedentary population Tomboctou and Segou	5-7/84	0-4		2000	0.10		0.20							Z	1022			
					734			3.54											
		1985	children																
		0-5.99	N.S.							0.50									
											0.00								
		0-5.99	6-10							0.50									
											0.00								
		4/86	0-5.99		195	7.64			0.00	0.00			0.51	0.51	8.15	B	7015		
					104	10.58			0.96	0.00			1.92	0.00	11.54				
		0-5.99	6-10		159	5.04			0.00	0.00			0.63	0.00	5.04				
					96	8.33			0.00	0.00			0.00	0.00	8.33				
		0-5.99	6-10		884	6.41			0.00	0.00			0.34	0.00	6.41				
	337			9.50			0.00	0.00			1.48	0.00	9.50						
0-5.99	N.S.							0.00	0.00		0.40	0.08	6.53						
									0.00	0.00		0.32	0.00	6.51					
6-10	F							0.00	0.00		0.49	0.16	6.38						
									0.19	0.00		1.30	0.00	9.67					
6-10	F							0.00	0.00		0.37	0.00	7.84						
									0.37	0.00		2.25	0.00	11.61					
1989	Mourdiab: Nara	5-14	B		478	6.69													
					208	2.90													
					270	9.60													
1990	Douen Kolokani (NW Bamako)	4-7	B			9.00		2.40			1.80			Z	7001				
					162	4.90		1.20		1.80									
MAURITANIA	USAID survey Adar Nouakchott	1983	<5	B	N.S.								2.60	Z	1042				
		1984	2-7										1.18		7001				
		1989	7-10										0.12		1042				
		1990				3118							0.64						

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.			
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *					
MOZAMBIQUE	Zambezia Province	1987-88	children	B	3790	7.30					0.25	0.89				7001			
	Maputo, Beira, Niampula	1990	5-6.00		10 267	0.30		0.30				0.10		0.70		1075			
NIGER	Tahoua, Maradi, and Zinder	5/1986	0-5.99	B	1388	3.97		0.72	0.07				0.22	0.14	4.83	7015			
				F	N.S.	3.77		0.72	0.14				0.14	0.14	4.63				
				M		4.18		0.73	0.00				0.29	0.15	5.06				
				B	6-10	611	4.75		0.65	0.00				0.82	0.00	5.40			
				F			5.52		0.00	0.00				0.58	0.00	5.52			
				M			3.76		1.50	0.00				1.13	0.00	5.26			
				B	0-5	425	7.33		1.18	0.24				0.24	0.24	8.75			
				B	6-10	179	7.82		0.56	0.00				0.56	0.00	8.38			
				B	0-5	386	3.11		0.52	0.00				0.52	0.26	3.89			
				B	6-10	213	5.16		0.94	0.00				0.47	0.00	6.10			
				B	0-5	422	3.21		0.47	0.00				0.00	0.00	3.68			
				B	6-10	169	1.78		0.00	0.00				1.78	0.00	1.78			
				B	0-5	1233	3.08		0.73	0.08				0.24	0.16	3.97	B		
				B	6-10	561	4.99		0.53	0.00				0.89	0.00	5.52			
				B	0-5	155	0.00		0.65	0.00				0.00	0.00	0.65			
				B	6-10	50	2.00		2.00	0.00				0.00	0.00	4.00			
					Gourc	1986	0-5		430	1.20									
	National	1988	0-6		1504	2.01							0.81		7004				
	Tillaberi Department	1990	0.5-6		2960	3.50	0.30	0.50				0.00	0.00						
			0-99		243	0.80													
			1-1.99		385	1.30													
			2-2.99		475	2.90													
			3-3.99		437	4.30													
			4-4.99		343	6.70													
			5-5.99		367	6.00													
			0-5		703	1.80										Z	7016		
	- Filingue		444	0.50															
	- Kollo		481	6.90															
	- Ouallun		297	0.30															
	- Say		666	6.50															
	- Tern		369	3.00															
	- Tillaberi																		
NIGERIA	National	1994	0-5.99	B	2836	1.00		0.15	0.60	0.00	0.00	0.20			Z	1061			
RWANDA	Sub-National - Gikongoro Prefecture	1987	0-6	B	5687			1.30				0.10	0.02		Z	7016			
					Gikongoro:	454		0.88											
					Karuma	257		2.33											
					Karunbo	424		1.41											
					Kinyamakar	556		2.88											
					Kivu	438		1.14											
					Mubagu	710		0.84											
					Mudasomwa	608		1.97											
					Muko	415		0.96											
					Musange	701		0.85											
					Nshili	332		0.60											
					Nyamagabe	461		0.65											
Rukondo	331		1.21																
Rwumikowa																			
SENEGAL	Casmance	11-12/79	all	B	1262							0.08			Z	7012			
					1988	865	14.10	2.70	0.20						7016				
					Diourbel and Fatick - peanut growing														
	Maficounda	2/1990	2-14		1259								0.00		1072				
					2/1991	1008							0.00						
TOGO	Kara: Bassar	1992	0-4	B	N.S.	10.00									Z	7016			
					Savanes: Dapaong		10.00												
UGANDA	Kamuli District	1991	0-6.00	B	5074	2.70			1.00			1.70	0.30	4.00	B	7004			
					- preliminary results														
					- dry season	10-11/91	0-5	5003	2.64		1.14				1.70	0.30	5.38		
							0-0.99	942									1.60	Z	7023
							1-1.99	765									1.20		1032
							2-2.99	773									2.30		
							3-3.99	871									4.00		
							4-4.99	751									5.90		
							5-5.99	901									8.40		
						0-5	1325										6.60		
	- Budiope																		
	- dry season																		
	- Bugabula																		
	- dry season																		

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' - Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms								Total *	Ref. No.	
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal			
UNITED REP. OF TANZANIA	- Bulamogi dry season				830									4.00		
	- Buzaaya dry season				1779									2.00		
	Iringa	1974	0-4.99	B	N.S.		0.15	0.15	0.51			1.17	0.56	0.86	C	1024
	Mbeya, Iringa & Kagera	1982-85	0-5.99			12 980		0.21	0.10			0.85			Z	1014
			0-99			2299		0.00	0.00			0.13				
				1		2999		0.13	0.03			0.20				
				2		3051		0.07	0.07			0.39				
				3		2675		0.34	0.04			0.41				
				4-5		1956		0.36	0.46			0.77				
	National - 11 Regions	1983-85	0-4			15 593		0.14	0.12	0.05		0.31				1025
	Biharamulo					2661		0.41	0.11	0.00		0.04				
	Iringa					1760		0.11	0.06	0.40		0.63				
	Igungu					735		0.00	0.54	0.00		0.14				
	Ludewa					1881		0.00	0.00	0.00		1.12				
	Mafinga					1180		0.25	0.00	0.00		1.27				
	Mbeya					188		0.00	0.00	0.00		0.00				
	Ngara					1776		0.34	0.11	0.00		0.00				
	Njombe					881		0.11	0.23	0.34		1.02				
	Nzega					2621		0.00	0.27	0.00		0.04				
	Tabora					1235		0.00	0.00	0.00		0.00				
	Urambo					675		0.00	0.00	0.00		0.00				
	8 regions	1984	0-10			7431	0.30		0.37				0.86	1.53	B	1024
	Arusha					1418	0.60		0.10				1.10	1.80		
	Bukoba					623	0.20		0.20				1.60	2.00		
	Dodoma					1690	0.30		0.90				0.80	2.00		
	Iringa					124	0.00		1.60				2.40	4.00		
	Mbeya					1317	0.00		0.00				0.00	0.00		
	Morogoro					233	0.00		0.00				3.00	3.00		
	Singida					908	0.00		0.20				1.30	1.50		
	Tanga					1118	0.40		0.50				0.20	1.10		
	Moshi					8106	0.20		0.20				0.90	1.30		
	Iringa	03-06/84	0-4			5975		0.08	0.07	0.12		1.15				1014
	11/84				1331		0.23	0.23	0.00		2.70					
Tabora	03/85				5266		0.02	0.21			0.34					
				F	2501		0.04	0.00			0.24					
				M	2765		0.00	0.40			0.43					
Iringa	04/85			B	1049		0.00	0.19	0.00		0.38					
Kagera	07/85				4437		0.41	0.11	0.00		0.02					
Lusu Ward: 4 villages in Nzega	2/86	children			2380			0.67							1027	
Tabora	02/86	0-4			3177			0.60			0.47				1014	
				F	1522			0.39			0.46					
				M	1655			0.79			0.48					
ZAMBIA	3 districts: Luapula Province	8/85	0-5.99	B	4275	1.26		0.44	0.02		0.68		1.89	Z	1034	
			0-0.49		373			0.00			0.00	0.00	0.00	D		
			0.5-0.99		347			0.00			0.58	0.00	0.00			
			1-1.99		782	1.17		0.26			0.26	0.13	1.56	B		
			2-3.99		1466	0.76		0.27			1.02	0.00	1.03			
			4-5.99		1307	1.86		0.99			0.77					
			0-5.99		647	1.84					0.62		1.86	Z		
Kawambwa District					1086	0.71		0.09			0.55		1.11			
Mwense District					2542	1.36		0.75			0.71	0.04	2.24			
Nchelenge District																
Ndola:	1988	0-4.99			423	1.40		0.70							7016	
Kansensi					133	0.00		0.00								
Masala					97	1.00		2.10								
Chipulukusu					113	0.90		0.00								
Ndola - rural					80	5.00		1.30								
Ndola - schools:			6-12		1164	1.89	0.00	1.80								
Chibolele					309	2.00		1.20	0.00		0.00					
Kabwata					146	6.50		3.40	0.00		0.00					
Kansensi					232	0.30		0.30	0.00		0.00					
Lumano					79	5.10		1.20	0.00		0.00					
Ndola: Masala					398	1.20		2.40	0.00		0.00					
ZIMBABWE	Matabeleland- 6 districts	1991	children	B	6944	0.58		0.14				0.23		Z	1016	
Binga					1604	1.60		0.00				0.56				
Bubi					626	0.00		0.32				0.00				
Hwange					1249	0.56		0.40				0.00				
Lupana					882	0.00		0.00				0.00				
Nkayi					883	0.40		0.11				0.00				
Tsholotsho					1700	0.24		0.12				0.40				

* Total xerophthalmia rate: 'A' - XN-X3B, 'B' - XN-X3B, no X1A, 'C' - X1A-X3B, 'D' - X1B-X3B, 'Z' - Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.		
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *				
REGION OF THE AMERICAS																		
BOLIVIA	National - (INAN)	1981	5-4.99	B	5745	1.10								Z	2016			
	National - (PDRD)	1985	1-5		1088	2.30												
	La Paz :	1986	0-4		1969	5.00												
	Iturralde																	
	Inquisivi	1987	1-5		972	1.00												
BRAZIL	Coastal - total harvest	1981-1982	0-12	B	1440	0.00		0.07	0.00			0.00			2009			
	Coastal Bayeux				689	0.00		0.00	0.00			0.00						
	Coastal Mamanguape				751	0.00		0.13	0.00			0.00						
	- harvest																	
	- inter-harvest		0-4		1011	0.00		0.00	0.00			0.00						
	Sertao - Conceicao		0-12		1831	0.16		0.71	0.06			0.06						
	harvest																	
	- Itaporanga		0-4		917	0.22		0.44	0.00			0.00	0.00					
	harvest																	
	- Itaporanga		school		649	0.15		0.31	0.00			0.15						
	inter-harvest																	
	- Itapuranga		0-12		1566	0.13		0.38	0.00			0.06						
	harvest																	
	- Pianco - harvest				1594	0.25		0.63	0.00			0.19						
	- Total harvest				4991	0.18		0.58	0.02			0.11						
				0-5.99		2802	0.18		0.28	0.00			0.11					
				6-12		2189	0.18		0.96	0.05		0.09						
	Transitional Esperanca		0-4		770	0.00		0.00	0.00				0.00					
	- harvest																	
Transitional Esperanca				720	0.00		0.13	0.00					0.00					
- inter-harvest																		
Transitional-Esperanca			0-11		1431	0.00		0.00	0.00		0.00							
- harvest																		
Paraiba	1981-83	0-4.99		12 323	0.06		0.50				0.03			2002				
- Semi-arid				7070	0.07		0.78				0.06							
- Litoral				1762	0.00		0.06				0.00							
- Transitional				3491	0.06		0.17				0.00							
Parelhas Jardim de Serito	1986	0-6		6291	0.01		0.60							2024				
DOMINICAN REPUBLIC	Santa Domingo - hospital studies	1991	N.S.	B	820							2.80	Z	7010				
EL SALVADOR	National	4/73	1-6	B	9508			0.05			0.03		Z	7011				
GUATEMALA	Nebaj - Quiche	05/84	0-5	B	576		0.69				1.39			2004				
	Rural	1986	0-10		1369							0.50	Z	7010				
HAITI	National	1975	children	B	5589						2.50	8.00	Z	7002				
SOUTH-EAST ASIA REGION																		
BANGLADESH	Matlab - rural	1981	all	B	182 976	0.18		0.18				0.02	0.01	0.37	B	5032		
				F	90 564	0.10		0.10				0.02	0.01	0.21				
				M	92 412	0.26		0.26				0.02	0.01	0.53				
				0-6.99	B	39 120	0.35		0.36				0.02	0.04	0.75			
					F	18 735	0.25		0.26				0.03	0.04	0.55			
				7-19.99	M	20 385	0.44		0.46				0.01	0.03	0.93			
					B	61 258	0.29		0.29				0.04	0.01	0.59			
					F	30 335	0.13		0.13				0.03	0.010	0.27			
					M	30 923	0.45		0.45				0.05	0.00	0.90			
				National	1982-83	25-5.99	B	22 335	3.48	2.03	1.02	0.04		0.27	0.11	4.61		5057
								18 660	3.60	2.00	0.90	0.04	0.25	0.10	4.62			
				- rural				3675	2.80	2.50	1.60	0.05	0.35	0.16	4.54			
				Ranjpure District	1986	0-8.99		3040	3.55									5084
- baseline survey																		
- post intervention	1989			3389								1.74						
				F	1610							1.68						
				M	1779							1.80						
BHUTAN	National	1976	6-12	B	953							1.30	Z	5034				
	Central Zone		all		200							2.50		5033				
	Eastern Zone		6-12		151							2.20		5034				
			all		201							1.50		5033				
	Southern Zone		6-12		380							1.00		5034				
			all		484							1.00		5033				
	Western Zone		6-12		422						1.20			5034				

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *		
INDIA	National	1989	all		475									1.10	5033	
			0-5		3273									0.70		
	Coimbatore District	N.S.	0-6	B	3907			9.80					0.07	Z	5058	
	Assam		1-12		5039	5.22			0.04	0.06	0.00	0.15	0.10	11.27	A	5007
	- rural				2871	4.80			0.00	0.03	0.00	0.14	0.03	9.50		
	- tea garden colonies				1068	9.92			0.09	0.19	0.00	0.37	0.28	19.20		
	- urban				1100	1.81			0.90	0.00	0.00	0.00	0.90	8.27		
	7 villages in Madhya Pradesh: Jabalpur	< 1978	0-2.99		1000		24.10	7.50	1.10		0.20		1.30	32.90	C	5041
	9 states		1-5		N.S.									4.60	Z	
	Gujarat		pregnant school	F										26.00		5045
				B		16.00								41.00		
	Orissa		1-5											12.80		5041
	Baroda City		5-15	M	1704	6.00	18.00	2.00						31.00		5044
	- underprivileged															
	- one school		7-15	B	378	12.00	23.00	4.00						32.00		
	National - 10 states	1975	1-4.99		N.S.				0.60							5042
			5-12						1.70							
	NNMB Survey	1975-79	1-5						2.00							5043
	National - 10 states	1976	1-4.99						1.40							5042
			5-12						4.10							
		1977-78	1-4.99						1.40							
			5-12						4.20							
	9 States - Other villages	1977	0-4.99					13.30	4.90				0.60	18.80	C	5041
	- Tribal villages							12.60	4.20				0.20	17.00		
	Andhra Pradesh		1-5											4.40	Z	
	Gujarat													1.40		
	Karnataka													3.60		
	Kerala													0.70		
	Madhya Pradesh													8.70		
	Tamil Nadu													6.30		
	Uttar Pradesh													6.00		
	West Bengal													1.00		
	9 States - urban slums	1979	0-4.99				17.60	1.80					1.20	20.60	C	
	National - 10 states		1-4.99						0.90							5042
			5-12						1.30							
		1980	1-4.99						1.50							
			5-12						3.10							
		1981	1-4.99						2.70							
			5-12						5.10							
	Baroda City - underprivileged		5-15				16.00	30.00	8.00					41.00		
				F										37.00	Z	
				M										45.00		
			5-9.99	B										33.00		
			10-15											49.00		5044
	Sevagram	1981-82	0-99											0.00		5040
		0-5.99			1180	5.80	10.20	3.80					9.60			
			F		560	7.10	9.50	4.10					18.20			
			M		620	4.50	10.90	3.50					15.60			
		1-1.99	B		209								14.40			
			F		96								13.50			
			M		111								15.30			
		2-2.99	B		281								15.00			
			F		117								20.60			
			M		164								11.00			
		3-3.99	B		192								39.50			
			F		87								43.60			
			M		105								36.10			
		4-4.99			82								54.80			
		4-4.99	F		92								44.50			
		0-4.99	B		N.S.			1.80							5042	
Andhra Pradesh	1982							1.80								
Gujarat								1.80								
Karnataka								1.60								
Kerala								0.00								
Maharashtra								3.10								
Orissa								0.80								
Tamil Nadu								3.80								
West Bengal								4.90								
Gorakhpur	11/84				3710	2.35	0.65	0.86	0.08			0.46	0.08	3.75	5007	
Jodhpur					1877	3.62	8.84	6.82	0.05			0.80	0.10	11.29		
Ranchi					3959	4.98	8.26	2.98	0.03			0.28	0.13	8.17		
Calcutta					3905	4.05	5.99	3.12	0.18			0.18	0.05			
Baroda Survey - Chandrapur and Panchmahal	1986-87	0-6			4032	0.50	5.70	3.30					0.50		5085	

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *		
			0-1		N.S.	0.10	0.90	0.70					1.01			
			1-2			0.50	5.00	1.20								
			2-3			0.70	5.90	3.90					0.70			
			3-4			0.70	8.80	3.60					0.40			
			4-5			0.60	6.80	5.40					0.50			
			5-6			0.40	8.10	7.30					1.50			
	Kashi Block, Varanasi	1987	0-5		2304	6.80	12.10	8.20	0.20			0.70	0.54	15.90	A 5038	
	- total				1406	7.70	13.10	9.00	0.10			0.90	0.60	17.7		
	- PEM1&2				239	9.60	18.00	10.00	1.70			0.40	2.50	21.7		
	- PEM3&4				1706	1.40	9.00	0.90	0.00			0.50	0.00	11.30		
	- no morbidity				232	29.30	15.90	36.30				0.90	4.30	85.80		
	- with diarrhea				24	33.30	8.30	50.00					12.50	99.98		
	- ARI				260	25.40	28.10	34.60				1.20	1.60	89.70		
	- with worms				659	3.90	7.80	5.60	0.00				0.00	17.30		
	- normal nutrit. status															
	Gorakhpur - slums		0-6		1376	0.58	9.18	1.70	0.29			0.07	0.36	11.82	A 5036	
	- initial survey				1379	0.29	7.21	1.20	0.14			0.07	0.21	8.91		
	- 6 months later				15 775										5037	
	Andhra Pradesh, 5 health centers	1987-89	1-5													
	- baseline				7691			1.30								
	- after treatment				8084			2.90								
	- control							0.70							5043	
	NNMB survey	1988-90	0-5		N.S.										7003	
	Southern India	1989	0-4.99											11.00		
	Tamil Nadu- drought prone		6-4.99		15 419	3.70		7.20				0.07	0.05	10.95	B 5035	
	Chandrapur: Maharashtra	3-7/91	0-6		N.S.									0.99	Z 5064	
	Panchmahal: Gujarat													3.50		
INDONESIA	Central Java - plateau	07-08/73	1-4.99	B	2812		3.06	4.23							Z 5074	
	Central Java - plateau				1374		5.17	4.80								
	- rural				1438		1.04	3.69								
	- urban				360		1.00								5046	
	Medan town	11/74	0-5	F			1.10									
	4 villages - N. Sumatra		0-6	B	1017		16.80	0.70				1.10	0.40	17.90	C	
				F	489		11.50	0.60				1.00	0.60	12.70		
				M	528		21.70	0.80				1.10	0.20	22.70		
			0-0.99	B	207		5.80	0.00				0.50	1.00	6.80		
				F	108		3.70	0.00				0.90	0.90	4.60		
				M	91		8.80	0.00				0.00	1.10	9.90	C	
			1-3.99	B	328		13.40	0.30				1.20	0.60	14.30		
				F	158		8.89	0.60				1.30	1.30	10.80		
				M	170		17.60	0.00				1.18	0.00	17.64		
			4-5.99	B	490		23.50	1.29				1.22	0.60	24.70		
				F	223		17.00	0.90				0.90	0.00	17.90	C	
				M	267		28.80	1.50				1.50	0.00	30.30		
	National	1977-78	0-5.99	B	35 274			1.01				0.13	0.06		5017	
				I	5677			0.44								
				2	5857			1.20								
				3	5972			1.57								
				4	5475			1.28								
				5	6027			1.53								
	Aceh	1977	0-4.99		N.S.			2.42					0.48	2.90	D 5047	
	Bali							0.76					0.08	0.84		
	Bengkulu							0.65					0.22	0.87		
	Central Java							1.11					0.04	1.15		
	Central Kalimantan							1.50					0.11	1.61		
	Central Sulawesi							1.05					0.00	1.05		
	East Java							0.75					0.00	0.75		
	East Kalimantan							0.00					0.00	0.00		
	Jakarta							0.73					0.00	0.73		
	Jambi							0.91					0.00	0.91		
	Lampung							0.20					0.00	0.20		
	Moluku							2.04					0.00	0.00	Z	
	North Sulawesi							0.40					0.00	0.40	D	
	North Sumatra							0.43					0.00	0.43		
	Riau							0.43					0.00	0.43		
	South Kalimantan							1.47					0.00	1.47		
	South Sulawesi							0.40					0.08	0.48		
	South Sumatra							0.33					0.17	1.65		
	Southeast Sulawesi							0.49					0.00	0.00	Z	
	West Java							1.55					0.10	1.65	D	
	West Kalimantan							0.43					0.00	0.43		
	West Nusa Tenggara							1.57					0.21	1.78		

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.						
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *								
INDONESIA	West Sumatra	1983	0.5-5			2.80		1.31					0.16	1.47	5048							
	Lombok - pre program															0.90	1.60	0.22	0.22			
	Lombok - post program															0.90	0.24	0.20	0.04			
	Aceh - Baseline data	1982-83	0.5-99			24 986	1.17		1.23							5021						
																		0.00	0.10			
																	1	3867	0.29	0.44		
																	2	4148	1.35	1.16		
																	3	4294	1.82	1.84		
	4	3617	1.74	1.80																		
	5	5203	1.61	1.83																		
	West Java	1984					N.S.						0.00	0.14	5047							
	Bogor	1985				2169		0.60	0.83							5016						
																	- control-baseline					
																	- study-baseline	2102	0.38	1.24		
																	- control-6 month	2169	0.78	0.92		
																	- study-6 month	2102	0.33	0.33		
																	- control-12 month	2169	0.65	1.11		
	- study-12 month	2102	0.29	0.19																		
	Central Java						N.S.						0.00	0.91	5047							
	South Sulawesi	1986											0.00	0.26								
	West Nusa Tenggara												0.04	0.28								
	Aceh	1989	N.S.										0.10	1.38	7008							
	Central Java												0.00	0.91								
	Nusa Tenggara												0.04	0.28								
	South Sulawesi												0.00	0.26								
	4 Provinces:	1991	0-4			21 660								0.03		5047						
																	East Nusa Tenggara	5163	0.00	0.00	0.00	
East Timor																	5163	0.27	0.14	0.27		
Irian Jaya																	4339	0.25	0.00	0.00		
Maluku																	6798	0.10	0.00	0.00		
MYANMAR	Natogyi (townships)	1978-90	0-4.99	B	13 771			0.73							Z	5063						
- Saku					150			3.33														
- Thazi					5356			0.93														
- Wetlet					15 322			1.13														
6 townships	1982-85				49 263			1.00							5051							
4 regions:	1987-89	2-14			1862										5063							
																2-5.99	642	2.02				
																6-9.99	687	4.51				
																10-14	533	3.57				
																Hinthada	2-5.99	199	0.50			
																	6-9.99	196	1.53			
																	10-14	186	1.61			
																Kyauppadaung	2-5.99	185	4.32			
																	6-9.99	137	6.57			
																	10-14	158	3.80			
																Monywa	2-5.99	114	2.63			
																	6-9.99	223	8.52			
10-14	87	9.30																				
Taunggyi	2-5.99	144	0.69																			
	6-9.99	131	0.00																			
	10-14	102	1.96																			
Myin Gyan	1990	0-4.99			6030			1.21						5051								
Nyaung Oo					1917			1.62														
Pakokku					6551			0.50														
NEPAL	Baglung	1979	0-5	B	1489	1.00							1.00	Z	5068							
National	1979-80																					
																	7201	0.60	0.90			
																M	3632	0.70	1.10			
																F	3569	0.50	0.60			
																B	367	0.60	0.80			
																	- urban	6837	0.60	0.90		
																	- rural	2189	0.80	1.20		
																	- central	1833	0.80	0.90		
																	- eastern	147	0.30	0.50		
																	- far west A	790	0.60	0.90		
																	- far west B	3326	0.30	0.40		
																	- hills	258	0.00	0.00		
																	- mountains	3315	1.00	1.40		
																	- terai	305	0.00	0.00		
																	- valleys	1345	0.30	0.40		
																	- western	666	0.20	0.30		
																	Bagmati	622	0.00	0.10		
Bheri	91	0.00	0.00																			
Dhaulagiri																						

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.
						XN	X1A	X1B	X2	X3A	X3B	X5	Corneal	Total	*	
	Gandaki				561			0.30						0.70		
	Janakpur				686			0.70						0.70		
	Karnali				73			0.00						0.00		
	Koshi				669			0.30						0.50		
	Lumbini				693			0.30						0.30		
	Mahakali				450			0.80						1.30		
	Mechi				473			0.40						0.40		
	Naryani				837			1.50						2.20		
	Rapti				352			0.80						1.10		
	Sagarmatha				691			1.40						1.70		
	Seti				340			0.30						0.30		
	National	1981	0-14		6118	1.10		1.65				0.03	0.02			7005
			0-6		7580			0.64				0.20				
	Baglung	1985			478									1.70		5068
	Surkhet		0-4		1106	1.00										
	Lahun Eye Hospital	1986-88	0-10		4601				1.69			3.45		10.16	C	5086
				F	1793				1.56			4.41		10.32		
				M	2808				1.78			2.85		10.18		
			0-5	F	909				2.75			6.82		16.94		
				M	1285				3.50			4.82		17.89		
			6-10	F	884				0.34			1.92		3.50		
				M	1523				0.33			1.18		3.68		
	West Central Terai	1988	children	B	17 000									1.70	Z	7005
	Central Terai		0-10		8682			2.00								
	- Bara															
	- Parsa				8402			2.60								
	East Central Terai - Sarlahi		0.5-4		3895								0.05	3.05		
	Jumla	1990	0-4		3651									13.20		5059
	3 Terai Districts	1992	0-10		N.S.									4.10		5080
	FarWest Hill - Doti (20 clusters)	06-09/93	.5-5		1342				3.60							
	FarWest Terai - Kailali (20 clusters)				2116				2.20							
	FarWest Terai - Kanchanpur				1889				1.50							
	MidWest Mt. - Dolpa (20 clusters)				1245				4.30							
	MidWest Terai - Bardiya (20 clusters)				2206				3.30							
SRI LANKA	National	1975-76	.5-5.99	B	13 450			1.10							Z	5054
			.5-.99		1230			0.10								
			1		2598			0.20								
			2		2425	0.60		0.80								
			2-5.99		9536	1.00										
			3		2511	1.40		1.40								
			4		2503	0.90		1.40								
			5		2183	1.40		2.10								
	Amparni	1980-82	N.S.	F	N.S.	2.70		1.60			0.20	0.20				5062
				M		2.40		2.30			0.00	0.00				
	Anuradhapura			F		1.60		1.90			0.00	0.10				
				M		1.00		1.00			0.10	0.30				
	Badulla			F		0.40		0.40			0.00	0.10				
				M		0.70		1.10			0.00	0.00				
	Batticaloa			F		2.20		1.70			0.00	0.30				
				M		1.70		2.30			0.00	0.00				
	Columbo			F		0.30		0.70			0.00	0.00				
				M		0.10		0.80			0.00	0.10				
	Galle			F		5.20		8.70			0.00	0.20				
				M		2.60		7.60			0.00	0.00				
	Gampaha			F		1.10		1.20			0.30	0.10				
				M		0.60		1.70			0.30	0.00				
	Hambantota			F		0.50										
				M		1.40		2.30								
	Jaffna			F		0.60		2.50			0.00	0.50				
				M		0.70		2.90			0.10	0.20				
	Kalutara			F		2.10		0.90			0.00	0.20				
				M		1.50		0.40			0.00	0.20				
	Kandy			F		1.50		2.50			0.10	0.20				
				M		1.00		3.50			0.20	0.10				
	Kegalle			F		0.50		3.40			0.10	0.00				
				M		0.30		3.80			0.00	0.00				
	Kurunggala			F		0.20		0.20			0.00	0.00				
				M		0.20		0.80			0.10	0.10				
	Manar			F		4.10		0.40			1.80	0.40				
				M		1.80		0.60			1.00	0.40				

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.	
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *			
SRI LANKA	Matale			F		0.20		0.00			0.20	0.00					
				M		0.00		0.50			0.00	0.00					
	Matara			F		0.30		0.30									
					M			1.90									
	Moneragala			F				0.50			0.00	0.30					
					M			0.00			0.00	0.20					
	Mullaitivu			F		1.10		0.20			0.00	0.00					
					M		0.50		0.90		0.00	0.20					
	Nuwara Eliya			F					0.20		0.00	0.20					
					M		0.20		0.50		0.00	0.00					
	Polonnaruwa			F		0.70		0.60			0.00	0.00					
					M		0.70		0.80		0.10	0.10					
	Puttalam			F		0.00		0.00			0.00	0.00					
					M		0.00		0.30		0.00	0.00					
	Ratnapura			F		0.90		0.30			0.10	0.00					
					M		0.80		0.80		0.00	0.10					
	Trincomalee			F		2.20		5.10			0.20	0.00					
					M		1.50		5.10		0.30	0.10					
	Vavuniya			F					0.00		0.00	0.00					
					M				0.00		0.00	0.00					
	Columbo	1981	school	B	3973		0.40	0.20									
		1982			7198		8.30	1.20									
		1983			5522		2.40	0.20									
		1984			2546		6.50	0.00									
		1985			4210		5.20	0.10									
	National	1987	0-4		32 643	0.26	0.94	0.33									5026
			school		25 772	1.11	0.49	1.23									
Anuradhapura		0-4		3008	0.00	0.37	0.60										
Jaffna				6700	0.06	0.06	0.00										
Kulutara				4330	0.53	0.92	0.09										
Kandy				8917	0.37	0.79	0.07										
Kegalle				1245	0.08	2.33	0.64										
Kurunegala				5545	0.41	2.33	0.81										
Matale				1655	0.00	1.45	1.09										
Puttalam				1243	0.00	0.00	0.64										
THAILAND	Sakon Nakhon	05-06/85	3-7.99	B	271	0.00	3.90	0.00									
	- urban		3-3.99		51	0.00	6.50	0.00							Z	5029	
			4-4.99		56	0.00	0.00	0.00									
			5-5.99		37	0.00	6.40	0.00									
			6-6.99		58	0.00											
			7-7.99		69	0.00											
	- rural		1-7.99		1373	1.30	16.20	0.40									
			1-1.99		82	1.20	5.30	0.00									
			2-2.99		77	2.60	6.60	0.00									
			3-3.99		204	1.00	9.20	0.50									
			4-4.99		237	1.30	14.30	0.40									
			5-5.99		206	1.00	24.70	0.50									
			6-6.99		275	2.20											
			7-7.99		292	0.70											
	Sakon Nakhon, NE	1987	2-6		806	1.30		0.40								7005	
	North and Northeast - dry season	1990			996			1.00									
	North and Northeast Thailand	2-3/90			491	3.10										5070	
	South - 5 provinces	9-10/90			N.S.	2.20											
		1992			444						0.900	0.670				7005	
EASTERN MEDITERRANEAN REGION																	
AFGHANISTAN	Kabul	1989	5-14	B	N.S.		2.40								Z	3017	
			< 5				0.00										
DJIBOUTI	Rural	2/88	0-5	B	385	0.26		1.04			0.80				Z	3013	
	Urban				235	0.00		0.43			0.43						
IRAQ	Hartha - rapid assessment	05/91	1-6	B	231	2.90		1.40							Z	3021	
	Bughdad, Basrah, Mosul: health centres - (preliminary data only)	1994	0-4.99		7000	1.60								2.10	A	3026	
MOROCCO	National	1971	0-3.99	B	6710									1.00	Z	3008	
OMAN	12 MCH centres	1981	0-6	B	566		5.30	1.50							Z	3019	
				F	172		1.10	1.10									
				M	394		7.10	1.70									
	N.S.	1991	0-2	B	N.S.									0.00		3005	

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.			
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total *					
PAKISTAN	National micronutrient survey	1976-77	0-4.99	B	N.S.			1.50							Z	3003			
	Karachi	1976-79	0-3		110	2.73	0.00												
				F	57	1.75	0.00												
				M	53	3.77	0.00												
				B	385	26.23	1.82												
				F	212	17.45	1.42												
	Orangi, Al Fateh	1987	5-15	B	106	22.00	3.80									3014			
M				173	36.99	2.31								3006					
Thar Desert - drought			N.S.	N.S.	7.00										3027				
Karachi - 3 urban slums	2-3/90	.5-4.99			578								0.00						
SUDAN	Red Sea: Kassala	1980	0-6		1555	13.70			0.06					2.70	Z	3018			
				M	814	15.60			0.12					3.56					
		F	741	11.61			0.00					1.75							
	Sinkat West: Red Sea Province		0-15	B	100	12.00	8.00				0.00	0.00	20.00	C					
	Wad Kowli refugee camp	1985	children		104	5.70	10.60					2.90	1.00	17.30		3011			
	Gebeit Elma'adin: Red Sea Province	5/85	0-15		64	4.70	0.00					0.00	0.00	4.70					
	Muhammad Qod: Red Sea Province				67	10.40	3.00					1.50	0.00	13.40					
	Omduran camp-Kordofan Province		0-15		124	12.10	3.20					0.00	0.00	15.30					
	Quirba Central		N.S.	F	14	28.57	14.29					0.00	0.00	42.86					
	- health workers																		
	- lactating with therapeutic feeding			lactating	17	11.76	5.88					23.53	0.00	17.64					
	- lactating, gen. population				46	4.35	17.39					2.17	0.00	21.74					
	- program, gen. population			pregnant	5	0.00	60.00					0.00	0.00	60.00					
	Wad Sherife camp	1985-86	0-15	B		100	9.00	5.00					2.00	1.00	15.00				
	- supplementary feeding																		
	- therapeutic feeding							115	13.00	4.30					0.00	0.00	17.30		
	- totals							215	11.20	4.70					0.90	0.50	16.40		
	Asemi: Abetchi Bergu Tribe - outside camp	6/85			28	10.71	3.57						3.57	0.00	14.28				
	- inside camp				43	9.30	0.00						6.980	0.00	9.30				
	Asemi camp-Dudjo Tribe				45	8.88	0.00						0.00	2.20	11.10				
	Asemi camp-nomad				42	7.14	0.00						0.00	0.00	7.14				
	Zagat Arabs																		
	Fau refugee camp				88	5.70	11.40						2.30	4.50	21.60				
	- supplementary feeding																		
	- therapeutic feeding				44	0.00	4.55						2.27	0.00	4.55				
	Blue Nile	1986-87	0-5			N.S.	1.60										3015		
	Gezira						0.40												
	Kassala						0.50												
	Khartoum						2.10												
N. Darfur						3.50													
N. Kordofan						5.60													
Nile						0.70													
Northern						0.50													
Red Sea						1.30													
S. Darfur						4.50													
S. Kordofan						1.60													
White Nile						1.10													
North Darfur:	1988			<6		1919	0.52		0.10					0.10				3010	
Forig							21	0.00		4.80					0.00	0.00	4.80	B	3022
Heilat Salih					434	0.20										Z			
Humaida					88	0.00													
Imtidad					103	0.00													
Kheir Wagid					295	0.30													
Oordi					126	0.00													
Rumalia					302	0.30													
Sunger, Duhl. Kurge					288	1.70													
Shag Zaroog					205	1.00													
Tosal					70	0.00													
Rural					1787	0.60			0.00					0.34	0.06	0.66	B		
Urban					103	0.00			0.00					0.97	0.00	0.00			
Khartoum & Gezera Province - rural		1988-90	0.75-6				30 000	3.30		3.30								7004	

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms								Total *	Ref. No.	
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal			
YEMEN	Derudah	10-12/89	0.5-6		158									0.00	Z	3024
	Halayb				218								0.00			
	N. Tokar		272											6.20		
	Red Sea Province		<6	2001	1.10		0.70					0.05		1.60		
				332										0.30		
				276										1.10		
				403										1.50		
				302										1.70		
	Rural Port Sudan	<6	281										3.20			
	Rural Red Sea		144										2.10			
	South Tokar		1207										2.50			
	Sinkat		259										0.00			
	Urban Port Sudan		156										6.40			
	- A		194										0.00			
	- B		197										1.50			
	- C		204										0.00			
	- D		199										0.00			
Urban Red Sea		794										0.40				
6 health centres	1981	0-5.99	B	322				0.00	0.00	0.00					3012	
		6-14		267	1.12	10.00	4.00				0.37					
Tihama region - 21 villages (preliminary)	5-6/92	1-6		2470	0.40		1.70				0.04	0.04	2.14	B	3020	
			F	1148	0.2		1.22				0.09	0.00	1.42			
			M	1322	0.53		2.11				0.00	0.08	2.72			
			B	447	0.00		1.6				0.22	0.00	1.60			
				500	0.40		1.40				0.00	0.20	2.00			
				468	0.43		2.13				0.00	0.00	2.56			
				611	0.82		2.94				0.00	0.00	3.76			
WESTERN PACIFIC REGION																
CAMBODIA	5 provinces	1993	1-6	B	10 107	6.30		0.60	0.03				0.07		Z	6020
	Takeo				2000	11.00	0.60	0.50	0.05				0.01			
	Rattanakiri	06/93			1607	2.05	0.50	0.93	0.06				0.00			
	Koh Kong	07/93			2150	5.30		0.74	0.00				0.05			
	Kompong Thom				2288	9.80		0.69	0.04				0.05			
Phnom Penh				2062	2.08		0.00	0.00				0.10				
COOK ISLANDS	High-risk mainut. areas	1989-92	5-6	B									0.00	Z	7022	
KIRIBATI	Bonriki village	1989	<7	B	150	12.00	18.00	4.00							Z	6019
	Tabiang village				80	24.00	9.00	11.00								
Tarawa and Abemama Atoll	230				16.00	15.00	7.00									
	National - 6 islands	9-10/89	0.5-5		4614	3.52		10.85	0.32	0.04		1.08	14.74	B	6013	
				F	2169	3.50		9.00	0.10	0.00		0.80	12.60			
				M	2445	3.60		12.50	0.50	0.10		1.40	16.60			
	Abemama Island			B	321	8.40		13.70	1.60	0.00		1.90	23.60			
	Eutaritari Island				640	0.60		2.80	0.00	0.00		0.00	3.40			
	Nonouti Island				492	7.70		10.80	0.60	0.00		2.40	19.10			
	South Tarawa Island				1894	3.20		10.80	0.30	0.10		1.10	14.40			
	Tabiteuea North Island				594	4.00		15.20	0.20	0.20		1.90	19.50			
	Abaiang Island				686	1.30		13.50	0.00	0.00		0.20	14.80			
	Kiritimati Island - 3 villages	1994	0-6		357								1.12		6023	
	Banana village				106		2.83									
	Tabakea				117		0.85									
	London				134		0.00									
LAO P.D.R.	Mekong Valley	1968-69	N.S.	B	2988								7.62	Z	7005	
	National	1995	2-5.99	B		0.70									6026	
MALAYSIA	Iban-Sut and Mujong River	1976-78	0-6	B	414									20.50	Z	6006
	Iban - Lemanak River				388									12.90		
	Iban - middle Mukah River				460									3.90		
	Kayan and Kenyah - Baram River				556									11.50		
	Land Dayak-Febakang				552									6.90		
	Malay - Sarawak River Delta				361									38.20		
	Melanau - Tillian River 3rd				352									2.00		

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms										Ref. No.		
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal	Total	*			
MARSHALL ISLANDS	Penan - Mulu area Sungai Choh	1977	0-4.99 adults school		131									19.10				
					651								1.60	B				
					202								5.50	Z				
					251								10.80					
MICRONESIA (FED. STATES OF)	National - Chuuk	12/88-5/89	3-6.99	B	448	14.00		6.00							Z	6017		
PAPUA NEW GUINEA	9 provinces - hospitals	4-6/1993	.5-5.99	B	1027	0.39		0.19					0.00	0.59	Z	7023		
PHILIPPINES	Cebu	1976	N.S.	B	N.S.									4.50	Z	7003		
	Metro Manila	1976-82	N.S.											3.90				
	Cebu and Marinduque	1979-81												4.40				
	National	1982	2-6		2621	1.80		1.40						3.50		6022		
		1987	5-6		3389	0.70		0.20				0.10	0.00	0.90				
	Antique		N.S.		N.S.	2.40		0.80						4.00				
	Las Pinas		N.S.		N.S.	1.20		2.10						3.60				
	Antique	1989	0-5.99		4948	0.80		0.20					0.04	1.04	B	7005		
	N. Samar	1991	N.S.		3127	4.40		2.70					0.20	7.30				
	Quezon		0-5.99		3404	1.60		0.60					0.00	2.20				
	Zamboanga		N.S.		4847	1.50		2.00					0.10	3.60				
	SOLOMON ISLANDS	National nutrition survey	1993	5-6	B	5049	0.40		0.10									
SOLOMON ISLANDS	Gizo Island Guadalcanal Kolombangara Malaita Marovo Lagoon Vella Lavella Vonavona Lagoon 7 islands	11-12/91	5-6	B	273									2.20	Z	6014		
					339								0.60					
					106								5.70					
					1495								1.50					
					246								1.20					
					342								0.30					
					95								5.30					
					2896	0.52		1.42				0.00	0.07	1.52	B			
														0.60	Z			
														0.80				
									1.30									
									3.90									
									20.80									
TUVALU	National	1991	5-6	B	1059	0.00		0.00					0.00	Z	6025			
VANUATU	National	1991	<5	B	1870								0.05	Z	7023			
VIET NAM	creches, kinderg, orphans, general population	1984	0-5	B	1700								0.65	4.20	Z	6010		
VIET NAM	National	1985-87	0-4.99		34 214	0.37		0.16				0.12	0.07	0.60	B	6016		
					5038	0.67		0.22				0.00	0.89		6011			
VIET NAM	National (totals)		0-4.99		21 337	0.29		0.12				0.12	0.07	0.48				
				F	10 734	0.18		0.08				0.14	0.06					
				M	10 643	0.39		0.16				0.10	0.08					
	Dac Lac		0-4.99	B	501	0.19		0.40				0.00	0.00	0.59				
	Ha Nam Ninh				3147	0.25		0.06				0.19	0.13	0.44				
	Ha Son Binh				1286	0.15		0.00				0.08	0.78	0.23				
	Hai Phong				557	0.00		0.00				0.00	0.00	0.00				
	Hanoi				7886	0.34		0.16				0.11	0.76	0.58				
	Kien Giang				749	0.27		0.40				0.00	0.00	0.67				
	Long an				651	0.31		0.00				0.00	0.15	0.46				
	Minh Hai				621	0.32		0.32				0.48	0.00	0.64				
	Nghe Tinh				1144	0.26		0.00				0.09	0.09	0.35				
	Thanh Hoa				3092	0.23		0.00				0.03	0.06	0.29				
	Vinh Phu				1723	0.46		0.23				0.29	0.06	0.75				
	Central Highlands	1985-89			2455	0.48		0.28				0.24	0.00	0.76		6016		
	Central Vietnam	1985-88			501	0.40		0.40				0.00	0.00	0.80		6011		
	- High Plateau																	
	- North				4236	0.43		0.00				0.02	0.07	0.52				
	- South Coast				2405	0.91		0.41				0.16	0.12	1.44				

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

Country	Geographic area	Survey year	Age group	Sex	Sample size	Prevalence of ocular signs and symptoms								Ref. No.		
						XN	X1A	X1B	X2	X3A	X3B	XS	Corneal		Total *	
	Hanoi City	1985-89			1180	0.17		0.00				0.17	0.00	0.17	6016	
	Ho Chi Minh City				2514	0.36		0.13				0.27	0.13	0.62		
	Mekong River Delta				2021	0.34		0.25				0.15	0.05	0.64		
	Midlands				1723	0.58		0.22				0.30	0.06	0.86		
	Mountains				5149	0.27		0.14				0.09	0.04	0.45		
	North of Central Coast				4236	0.43		0.00				0.02	0.07	0.50		
	Red River Delta				11 716	0.43		0.13				0.12	0.09	0.65		
	South of Central Coast				5734	0.78		0.31				0.08	0.07	1.16		
	National	10/88	0-4.99		23 782	0.35	0.15	0.15				0.13	0.07	0.57		A 6015
				F	11 930	0.27	0.13	0.10				0.13	0.07	0.57		
				M	11 852	0.44	0.17	0.20				0.12	0.09	0.90		
			0-0.99	B	3938								0.05	0.13		
			1-1.99		5470	0.05	0.00	0.00	0.00				0.09	0.43		
			2-2.99		4872	0.51							0.16	1.00		
			3-3.99		4844	0.60							0.08	1.13		
			4-4.99		4685	0.58								0.90		
		1994	<5	B	37 920	0.05		0.04				0.05	0.01	0.14	6027	

* Total xerophthalmia rate: 'A' = XN-X3B, 'B' = XN-X3B, no X1A, 'C' = X1A-X3B, 'D' = X1B-X3B, 'Z' = Not specified

**PREVALENCE OF SERUM RETINOL
LEVELS SUGGESTIVE OF A SIGNIFICANT
PUBLIC HEALTH PROBLEM**

- DISAGGREGATED TABLES -

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A												
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (umol/l)					Ref. no.	
						*	< 0.35	< 0.70	Mean	S.D.		
AFRICAN REGION												
BURKINA FASO	Upper Volta	03-04/78	0-5.99	B	17	Z	17.6	70.5	0.58	0.25	7012	
			6-14		178		7.3	66.2	0.61	0.23		
			pregnant		78		6.4	57.6	0.70	0.33		
CAMEROON	8 villages: Kaele Dept. (N. Province) Extreme northern province	1988	.25-1.00	B	N.S.	Z	50.0				7001	
			1-1.99					19.0			1049	
			2-2.99					40.0				
			3-3.99					19.0				
			4-4.99					11.0				
CAPE VERDE	San Vicente and Fogo	1983	7-12.99	D	254	Z	6.7				7016	
			13-19		247		7.3					
			Periurban		95		1.1	17.9				
			Flooded plains		52		1.9	23.1				
CONGO	Brazzaville - past history of malaria - no history of malaria - malaria at time of survey	1988	.5-6.99	B	68	Z	0.0	38.2	0.99	0.52	1070	
					52		1.9	15.4	1.10	0.50		
					48		37.5	83.3	0.52	0.33		
CÔTE D'IVOIRE	Dabakala, Boniérédougou & Foumbolo	11-12/88	1-3.99	B	151	A	25.0	68.0			1059	
			- malnourished		96		31.3	68.8				
	- well nourished		55		14.5	67.2						
	NW: Odienne, Touba, Seguela & Mankono - after supplementation (Preliminary study)	3-11/94	.5-4.99			342	Z	12.3	36.6			1063
						170			54.4			
						172			45.6			
						N.S.			44.0			
								43.7				
ETHIOPIA	National	1980-81	0.5-5.99	B	739	Z	16.4	59.6	0.62		1052	
				F	301				0.62			
				M	438				0.62			
			.5-9.99	F	22				0.62			
				M	17				0.62			
			1-1.99	F	35				0.59			
				M	42				0.69			
			2-2.99	F	41				0.71			
				M	46				0.66			
			3-3.99	F	55				0.56			
				M	85				0.63			
			4-4.99	F	65				0.59			
				M	110				0.62			
			5-5.99	F	83				0.59			
				M	138				0.62			
Cash crop zone		.5-6	B	91		20.9	69.2	0.53**				
Cropping zone				477		15.1	55.6	0.65**				
Ensete zone				36		16.6	77.8	0.50**				
Pastoral zone				138		18.1	62.3	0.62**				
Shoa Region				344				1.00**				
Mekaye village (rural) - Hararge Region		1990s	0-12.5		80		30.2	88.2	0.46**	1072		
GAMBIA	Keneba, Manduar & Kanton	01/88	2-5.99	B	157	A			0.41	0.021	1050	
		04/88			157				0.69	0.021		
		08/88			157				0.78	0.029		
GHANA	VAST Morbidity Study - Kassena Nankana	6/90- 8/91	.5-4.99	B	1455	A	15.8	73.4			1064	
					21 906		14.4	56.9				
	VAST Survival Study - Kassena Nankana	1991	0-6		607	Z	14.0	54.9			7016	
					9		11.1	33.3				
Upper East - Kassena Nankana					54		14.8	53.7				
					118		9.3	49.1				
					113		20.4	58.5				
					11		17.1	57.3				

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified
** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A												
Country	Geographic Area	Survey year	Age group	Sex	Sample size	*	Vitamin A Levels (umol/l)				Ref. no.	
							< 0.35	< 0.70	Mean	S.D.		
	Kintampo District - northern region - southern region	04/1994	4-5		122		13.9	52.2			1058	
			5-6		74		6.8	59.5				
			1-4,99		99	A		51.0				
					50		8.0	64.0	0.64			
				49		6.0	37.0	0.80				
LESOTHO	National Micronutrient Survey	1993	2-6	B	127	Z	13.4	78.0	0.55	0.20	1055	
MALI	South Mali - rainy season	07-08/78	0-5,99	B	16	C	12.5	75.0	0.58	0.26	7012	
			6-14		119		0.0	57.9	0.67	0.23		
			adults		253		2.7	30.0	0.85	0.28		
			pregnant	F	60		1.6	38.3	0.77	0.23		
	- dry season	03-04/79	0-5,99	B	57		14.0	71.0	0.58	0.23		
			6-14		197		0.5	45.6	0.74	0.25		
			adults		388		3.3	22.6	0.96	0.36		
			pregnant	F	92		4.3	34.7	0.85	0.36		
	Kolokani (Northwest Bamako)	1990	2-10	B	152	A	15.1		0.53	0.03	1030	
MAURITANIA	Hodh-el Gharbi: Kerkerat	1987	1-15	B	81	Z	10.0	60.6			1069	
	Hodh-el Gharbi: Limberha				125		0.0	22.5			7016	
NAMIBIA	17 villages	1992	2-6	B	290	A	3.1	20.4		0.89	0.28	1037 1068
NIGER	Refugee camps - nomads	1975?	< 5	B	16	A	12.5	37.5			1067	
			> 5		56		16.1	39.3				
			all		72		15.3	38.9				
			< 5		5		20.0	60.0				
	- sedentary		> 5		42		16.6	92.8				
			all		86		36.0	87.2				
			N.S.		39		58.9	84.6				
	Refugee camps - nomads - pregnant or lactating - sedentary - pregnant or lactating - nomads and sedentary-pregnant or lactating - total				F	27		18.5	63.0			
						14		50.0	71.4			
						41		29.3	65.8			
				B	199		27.1	65.3				
NIGERIA	Cross River State	1983?	birth	F	62	Z			0.85	0.08	1043	
				M	50				0.93	0.22		
			1.5 weeks	F	62				0.94	0.07		
				M	50				0.99	0.13		
			0.25	F	60				1.14	0.34		
				M	40				1.29	0.15		
			0.6	F	56				1.26	0.23		
				M	44				1.33	0.15		
			1	F	56				1.47	0.50		
				M	44				1.54	0.37		
			children	F	112				2.35	0.49		
				B	941				0.72			
SENEGAL	Casmance - South Senegal	11-12/79	0-5,99	B	51	C	5.8	45.0	0.70	0.25	7012	
			6-14		231		1.7	41.5	0.79	0.27		
			adults		620		0.6	10.6	1.09	0.39		
			pregnant	F	113		0.0	9.7	1.04	0.32		
	Diourhal, Fatick & Kaolack - normal by ICT - M+ by ICT - M- by ICT - deficient by ICT	04/88		0-4,99	B	92	A			0.45	0.04	1023
						23				0.43	0.09	
						45				0.46	0.06	
						13				0.31	0.08	
						221				0.52	0.03	
						185				0.43		
Gadiack & Diop N'doffenc	4/1989	6/1989	2-6	Z	185		20.9			1053		
			2-7	B	185		40.0		0.43	1071		
	Louga - Linguere Department	7/1991	2-4,00		271		7.4	71.5	0.61	0.22	1074	
SOUTH AFRICA	Besler Farm	11/91	3-6	B	N.S.	B			0.77	0.25	1044	
									0.71	0.20		
									0.71	0.20		
							169		5.0	49.0		0.73
UNITED REPUBLIC OF TANZANIA	Iringa	1982-85	children	B	23	Z	0.0	13.0			1028	
	Tabora				64		1.5	45.3				

* Laboratory Assessment 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'F' = Fluorescence, 'Z' = Not Specified

** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A											
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (umol/l)					Ref. no.
						*	< 0.35	< 0.70	Mean	S.D.	
ZAMBIA	Dar-es-Salaam - children with measles - children with no measles	1985			34				0.40	0.23	1027
	Lusu Ward: Nzeza District - children without xerophthalmia				13				0.89	0.21	
					551	A			0.69	0.05	
	Luapula	1985	0-5.99	B	N.S.	Z	16.5				7001
	Ndola:	1989	6-12								7016
	Kabwata				90				0.0	27.0	
	Kanshensi				98				0.0	3.1	
Lumano				18				0.0	33.3		
Masala				90				1.1	7.8		
Chibolele				110				0.0	20.0		
REGION OF THE AMERICAS											
BELIZE	National	1990	0-4.99	B	N.S.	A			10.0		7007
	National	1990s ?	2-8		494			0.2	6.1	1.03	2021
BOLIVIA	National	1991	1-5	B	891	A	0.1	11.3			2016
	Altiplano - urban				N.S.				7.0		
	- concentrated rural								19.3		
	- poor area - rural		children						17.6		7002
	- rural dispersed		1-5						13.7		2016
	Llano - urban								8.4		
	- concentrated rural								16.5		
	- rural dispersed								11.9		
	- poor area		children						12.9		7002
	Valle - urban		1-5						8.2		2016
- concentrated rural								9.3			
- rural dispersed								3.9			
La Paz			0-4.99					1.2	9.0		7007
BRAZIL	National (migrants)	1972	N.S.	B	1081	Z	8.7	25.3			2007
	Northeast				255		5.9	26.3			
	South				79		7.6	11.4			
	Southeast				671		9.8	26.3			
	West				73		9.6	27.4			
	Northeast		0-6		N.S.		4.9	22.7			7010
	Gamela and Agua Preta	1973 ?	pregnant	F	165		0.0	2.5			2007
	Northeast	1981	0-6	B	N.S.		6.4	23.1			7010
		1982					1.5	14.7			
	Paraiba	1982	0-4		328		2.1	15.8			2002
	- litoral				155		1.9	16.7			
	- semi-arid				107		3.7	20.5			
	- transitional				66		0.0	6.1			
	Northeast	1983	0-6		N.S.		3.3	17.5			7010
		1984	1-7					13.2			
	Sao Paulo		0-4				B	3.6	30.2		2008
	- urban slum study										
- baseline			2-8		276		1.8	48.8			
- after oral supplementation					99		2.0	86.8			
South - before supplementation					97		1.0	89.6			
- 20 days after suppl.					97		0.0	40.2			
Poor areas - 3 periurban communities					182		2.7	73.6			
South	1986	2-8		N.S.	Z	1.8	48.8			7010	
Bahia - semi arid	1989	0-6			563		15.3	54.7		2032	
CHILE	National	1960	0-15	B	N.S.	Z	3.1	21.6			7010
COLOMBIA	National	1960	0-15	B	N.S.	Z	1.2	16.0			7010
		1977	0-4.99				0.0	24.1			
			5-9				0.0	28.8			
	Northern Coast		N.S.					41.1			7020
COSTA RICA	National	1979	0-5	B	N.S.	Z	0.0	2.3			7010
		1981					0.0	1.8			
DOMINICAN REPUBLIC	National	1969	0-15	B	N.S.	Z		9.0			7009
	Southwest	1991	1-5		505		4.2	19.6			
			1-2		104		2.9	15.4			
			2-3		101		5.9	24.7			
			3-4		94		5.3	20.2			
			4-5		115		3.5	18.3			
		5-6		91		3.3	19.8				

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified

** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM												
VITAMIN A												
Country	Geographic Area	Survey year	Age group	Sex	Sample size	*	Vitamin A Levels (µmol/l)				Ref. no.	
							< 0.35	< 0.70	Mean	S.D.		
ECUADOR	- rural				N.S.		5.2	21.3				
	- urban						3.3	24.7				
	National	1986	0-4.99	B	1570	Z	0.2	14.1			7010	
	- rural							16.4				
	- urban							11.9				
	- Azuay	1993	1-4.99			345	A		25.6			2026
	- Chimborazo				255			17.7				
	- Cotopaxi				183			9.6				
	- Esmeraldas				220			18.1				
	- Manabi				559			14.8				
	- rural				573			21.9				
	- urban				386			12.9				
EL SALVADOR	National - INCAP data	1976	1-6	B	N.S.	Z		33.3			7011	
	National	1988	0-4					36.0			7007	
	- rural							40.8				
	- urban							32.7				
	4 rural communities - Ahuashapan	1980s ?	0-4		259			36.0			2015	
	Huatules				65		6.0	41.0				
	Magueyes				64		8.0	36.0				
Palo Pique				64		11.0	58.0					
Roble				66		3.0	26.0					
GUATEMALA	National	1970	0-4	B	N.S.	Z		26.2			7010	
	Guatemalan children	10-11/75	0-4		543	B	3.3	21.5	1.02	0.36	2010	
	- baseline sugar fort.											
	- 1 year sugar fort. trial	10-11/76			644		0.3	5.1				
	National	1976			0-5	Z		33.0			7010	
	Guatemalan children - 2 year sugar fortification trial	10-11/77	1-5		721	B	0.3	9.2			2010	
	National - rural	1988	1-4		N.S.	Z	3.2	21.6			7009	
	Western - HOPE/INCAP	1989	0-5					26.0			7010	
- high altitude	1991	<6		3250			20.0			7021		
- low altitude				N.S.			14.0					
								37.0				
HONDURAS	National	1987	0-4	B	N.S.	Z		20.0			7020	
MEXICO	Yucatan	1984	N.S.	B	N.S.	Z		25.9			7020	
	Hermosilla	1990	2-7		N.S.			32.0			7009	
NICARAGUA	USAID - preliminary results	1993	1-4.99	B	N.S.			7.9	31.3		7023	
PANAMA	National - 4 regions	1992	1-4.99	B	1103	B	0.0	6.0	1.31		2027	
	- indigenous				106		2.0	13.0	1.11	0.41	2033	
	- non-indigenous				976		0.0	5.0	1.33	0.42		
PERU	Lima hospital	1989	1.5-1.5	B	72	B			0.51	0.46	2034	
	- children with diarrhea											
	- healthy control				65				1.00	0.32		
	Piura- rural	1992	0-6		300	Z		32.8			7020	
	Puno- rural				N.S.			14.1				
UNITED STATES OF AMERICA	N-Hanes I	1971-74	3-5.99	B	689	A		2.0			2011	
						725			2.8			
						927			0.5			
						930			0.5			
		NHanes I	1971-74	4-5	F	485				1.27		2014
						501				1.28		
		- black				127				1.13		
						118				1.12		
		- white				354				1.27		
						381				1.31		
		National	1976-80			471	Z			1.16		
						493				1.13		
		NHanes - Mexican Americans				100	A			0.96		
						500				1.03		
		NHanes II	1976-80	4-5	F	418				1.10		
						500				1.16		
		- black				74				1.06		
						69				1.04		
	- white				335				1.10			
					413				1.18			
	New York City				75				1.26			
	- Pb screen healthy											

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified

** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM										
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (umol/l)				Ref. no.
						*	< 0.35	< 0.70	Mean	
	- measles and convalescent Puerto Ricans in NYC	1976-80	12-17	M	111				1.23	2011
	Mex-American SW, Cuban-American NHanes II				1026			0.0		
					1009			0.3		
			3-5.99	F	589			6.1		
				M	702			3.4		
		1982-83	6-11	F	604			2.3		
	Mexican Americans in SW				613			2.2		
			4-5.99	B	234			4.6	1.00	
				M	134			3.8		
				F	100			5.8		
			6-11.99	B	1028			2.7	1.11	
				F	521			2.9		
				M	507			2.4		
			12-17	B	982			0.1		
				F	496			0.2		
			M	486			0.0			
	Non-Hispanic	4-5.99	B	142			5.7	1.04		
	- blacks									
	- whites			654			2.5	1.15		
	- blacks	6-11.99		179			2.2	1.19		
	- whites			926			1.7	1.19		
SOUTH-EAST ASIA REGION										
BHUTAN	National	1985	0-4	B	134	Z	14.0			7008
	National - pregnant women		pregnant	F	149		13.0			
INDIA	Coinbatore district	N.S.	N.S.	B	20	Z		0.42	0.07	5058
	- control - initial				20			0.42	0.07	
	- control - 3 months				20			0.43	0.06	
	- control - 6 months				20			0.63	0.08	
	- control - 9 months				20			0.64	0.08	
	- control - final				20			0.40	0.11	
	- Papaya fed - initial				20			0.86	0.05	
	- Papaya-fed - 3 months				20			1.20	0.06	
	- Papaya-fed - 6 months				20			1.36	0.16	
	- Papaya-fed - 9 months				20			1.64	0.18	
	- Papaya-fed- final				20			0.43	0.10	
	- Amaranth-fed-initial				20			0.89	0.05	
	- Amaranth-fed-3 months				20			1.21	0.06	
	- Amaranth-fed-6 months				20			1.49	0.06	
	- Amaranth-fed-9 months				20			1.50	0.07	
	- Amaranth-fed -final				20			0.41	0.07	
	- Carrot-fed - initial				20			0.90	0.05	
	- Carrot-fed - 3 months				20			1.25	0.06	
	- Carrot-fed - 6 months				20			1.58	0.10	
	- Carrot-fed - 9 months				20			1.84	0.07	
	- Carrot-fed - final				20					
	Tamil Nadu - drought prone	1989	.5-4.99		280		21.4	37.5		5035
	Stun area in Gorakhpur		0-5		6	C		100.0	0.44	0.05
	- nightblind children									
	- XIA +				14		21.4	78.6	0.44	0.07
	- XIB +				8		25.0	75.0	0.42	0.09
INDONESIA	Nine regencies	1975-76		F	305	Z	17.0	51.0		5050
	- lactating				61		23.0	66.0		
	- pregnant				131		13.0	58.0		
	- representative			B	268		9.0	47.0		
	West Java (Putkakarta)	1978								7008
	Bandung - Bitot's spots	02-08/83	3.5-4.5		4	A			0.15	0.07
	- Bitot's spots 10days				4				0.70	0.20
	- Bitot's spots 21 days				4				0.42	0.12
	- corneal xerosis				5				0.27	0.18
	- corneal xerosis 10 day				5				1.12	0.73
	- corneal xerosis 21 day				5				0.84	0.45
	- nightblind				6				0.45	0.25
	- nightblind 10 days				6				0.52	0.29
	- nightblind 21 days				6				0.52	0.21
	- reduced growth				5				0.66	0.24
	- reduced growth 10 days				5				0.66	0.24
	- reduced growth 21 days				5				0.63	0.25
	- satis. growth 10 days				5				0.59	0.25
	- satis. growth 21 days				5				0.38	0.13
	- satisfactory growth				5				0.45	0.23
	South Sulawesi	1988	0-5		N.S.	Z	8.9			7005
	West Java						5.9			

* Laboratory Assessment: 'A' = HPLC, 'C' = Spectrophotometry, 'D' = Fluorescence, 'Z' = Not Specified

** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A												
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (µmol/l)				Ref. no.		
						*	< 0.35	< 0.70	Mean		S.D.	
	West Kalimantan	1991					15.6					
	Irian Jaya			484		13.6	68.8	0.60	0.28			
	Maluku			508		12.6	69.3	0.64	0.39			
	Nusa Tenggara Timur			536		9.1	58.0	0.71	0.37			
	Timor Timur			341		14.9	86.6	0.73	0.38			
	West Java (Bogor)			257	N.S.		1.2	32.7		7008		
	Rural Javanes community		0.5-0.99		96		10.0	69.0	0.59	0.18	5071	
				1-1.99		173		6.0	63.0	0.66	0.21	
				2-2.99		191		5.0	50.0	0.69	0.21	
				3-3.99		208		7.0	58.0	0.65	0.21	
				4-4.99		20		0.0	65.0	0.67	0.21	
	Cibungbulang Subdistrict : Bogor	11/12-92	15-40		73	A			30.1	0.98	0.44	5083
	West Java: Bogor - lactating women after oral dosing	1994	17-37	F	64					0.99	0.35	
West Java Bogor - non-pregnant control group after oral dosing		29-41		14					1.47	0.31		
MYANMAR	Kyaokpadaung	1987-89	2-14	B	480	Z	8.9	60.0			5063	
			2-5.99		185		16.1	76.1				
			6-9.99		137		11.1	49.5				
	Hinthada			10-14		158		1.3	53.6			
				2-14		581		7.9	18.2			
				2-5.99		199		11.0	22.9			
	Monywa			6-9.99		196		5.4	17.6			
				10-14		186		7.4	14.1			
				2-14		424		4.7	15.8			
	Taunggyi			2-5.99		114		2.3	19.5			
				6-9.99		223		7.6	18.3			
				10-14		87		2.5	7.5			
				2-14		377		0.0	6.7			
				2-5.99		144		0.0	10.8			
				6-9.99		131		0.0	3.5			
			10-14		102		0.0	4.9				
SRI LANKA	All children w/+ eye findings	10/75-3/76	.5-5.99	B	29	Z	7.0	21.0			5054	
	All children w/- eye findings				350		0.0	5.0				
	subsample - rural children				346			5.0				
				4-6		139			6.0			
	Ratnapura and Kandy		.5-6		178				1.12			
	- rural serum sample											
	- wasted only				16				1.20			
	- rural serum sample			4-6		63			1.17			
	- village serum sample			N.S.		141			1.14			
	- estate serum sample				37				1.06			
	- rural-wasted sample			4-6		5			1.13			
	- village-wasted sample			N.S.		10			1.29			
- estate-wasted sample				5				1.00				
Colombo - serum normal			4-6		137				1.35			
- special children					143			0.0				
- rural-wasted					2				1.19			
THAILAND	Ubon	1974-78	0-4	B	143	Z	17.0	87.0	0.53	0.03	5082	
			schoolage children	F	67		22.0	73.0	0.56	0.04		
	Ranathibodi hospital	1984	N.S.		B	43		2.0	44.0	0.81	0.04	
				- control group		15		7.0		0.84	0.10	
	- with measles				25		32.0		0.42	0.03		
	Sakon Nakhon		1-2.99			127		23.6	81.9			5029
				- rural								
					F	62		27.4	83.9	0.54	0.22	
					M	65		20.0	80.0	0.52	0.22	
				3-6	B	410		9.3	41.5			
					F	218		6.9	39.5	0.81	0.33	
					M	192		12.0	43.8	0.75	0.26	
				6-8.99	B	315		4.5	42.3			
					F	162		5.6	37.7	0.79	0.30	
					M	153		3.3	46.8	0.74	0.25	
- urban			3-6	B	100		0.0	21.9				
			3-6	M	48		0.0	20.8	0.90	0.23		
			3-6	F	52		0.0	23.1	0.87	0.22		
Sakon Nakhon	1986	3-9	B	50	A				0.60			
- baseline (supplemented)					45				0.54			
- control post intervention					50				0.75			
- supp. post intervention												
North and Northeast Province	1990	2-6			996		0.6				7005	

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' - Not Specified
 ** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A											
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (µmol/l)				Ref. no.	
						*	< 0.35	< 0.70	Mean		S.D.
	- dry season	02-03/90			485		0.6	14.9	1.02	0.34	5070
	Machongson				43			39.6	0.85	0.22	
	Mukdaharn				15			4.1	1.10	0.35	
	Nan				49			6.3	1.05	0.23	
	Payao				21			19.0	0.96	0.26	
	Pisanuloke				43			2.0	1.31	0.35	
	Sakonakorn				26			24.5	0.94	0.46	
	Surin				48			10.4	0.93	0.25	
	Khon Kaen	09-10/90			11			4.7	1.04	0.27	
	Machongson				43			3.8	1.62	0.25	
	Mukdaharn				24			2.1	1.89	0.81	
	Nan				54			24.1	1.07	0.23	
	Payao				26			10.9	1.14	0.35	
	Pisanuloke				53			2.0	1.61	0.35	
	Sakonakorn				49			5.0	1.27	0.42	
	Surin				49			3.9	1.26	0.35	
	North and Northeast Thailand - Totals				56			7.2	1.30	0.54	
	National Data	1991	6-11		11	N.S.	Z	3.0	39.2		5079
EUROPEAN REGION											
FRANCE	Tours	3/85-1/86	1-16	B	392	A			1.47	0.42	4005
				F	185				1.46	0.37	
				M	207				1.50	0.46	
			1-3	B	10				0.96	0.24	
			3-6		13				1.01	0.23	
			6-9		112				1.33	0.33	
			9-12		104				1.41	0.33	
			12-14		50				1.53	0.34	
			14-16		97				1.81	0.43	
ISRAEL	Arab children- low SES - E. Jerusalem	1970s ?	5.5	F	24	C	0.0	83.3	0.58	0.11	4001
				M	23		8.7	86.9	0.60	0.22	
ROMANIA	Orphanage	1970s	N.S.	B	117	Z			0.74	0.33	4003
TURKEY	Marmara, Central and East Anatolia	4/85-5/86	7-17	B	960	Z			1.18	0.46	4002
				F	60				1.24	0.55	
				M					1.29	0.58	
				F					1.42	0.77	
				M					1.18	0.61	
				F					1.32	0.48	
				M					1.34	0.76	
				F					1.23	0.56	
				M					1.28	0.55	
				F					1.42	0.70	
				M					1.32	0.73	
				F					1.34	0.75	
				M	30				1.90	0.35	
				F					1.19	0.47	
				M					1.57	1.35	
				F					1.26	0.73	
				M					1.26	0.51	
				F					1.29	0.54	
				M					1.27	0.52	
				F					1.22	0.37	
				M					1.07	0.36	

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified

** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A													
Country	Geographic Area	Survey year	Age group	Sex	Sample size	Vitamin A Levels (µmol/l)				Ref. no.			
						*	< 0.35	< 0.70	Mean		S.D.		
UZBEKISTAN	Small Towns City Villages Ankara - sick control group - measles group - health group	1994	17	F					1.37	0.66	4004		
			7-17	M	150				1.24	0.53			
				B	150				1.34	0.62			
			children		300				1.18	0.46			
					21			42.9	90.5	9.5		1.36	0.67
	Karakalpakistan - Muynak	5-6/1993	all	B	851	Z	16.4	48.9			4006		
	EASTERN MEDITERRANEAN REGION												
	DJIBOUTI	Djibouti - rural	1988	4-10	B	114	A	0.9	12.3			3013	
					F	N.S.			0.0	14.3			3028
					M				1.6	11.1			
				B	83			2.4	21.7			3013	
				F	N.S.			5.1	23.0			3028	
	M				0.0	22.0							
IRAN (ISLAMIC REPUBLIC OF)	4 provinces Kohgiluyeh and Boyer-Ahmad Tehran - goitre endemic area	N.S.	0-4.99	B	N.S.	Z		9.0			3016		
			6-12						6.4				
			13-17						2.7				
			7-17 children	F			0.0		0.0				
OMAN	National Muscat Dhofar Dhakiliya N. Sharqiya S. Sharqiya N. Batinah S. Batinah Dhahira	1995	5-6	B	759	A	2.1	20.8	0.94	0.34	3029		
				F	378			2.6	18.7	0.95		0.36	
				M	381			1.6	22.9	0.93		0.32	
				B	219			2.3	21.0	0.95		0.34	
				F	219			1.8	22.8	0.94		0.33	
				M	103			4.9	20.4	0.95		0.37	
				B	218			0.9	18.8	0.93		0.33	
				F	191			6.3	39.8	0.76		0.30	
				M	73			2.8	15.3	0.99		0.33	
				B	121			0.0	1.7	1.10		0.23	
				F	60			0.0	1.7	1.25		0.34	
				M	69			0.0	4.3	1.09		0.28	
				B	130			1.5	26.9	0.84		0.33	
				F	64			0.0	20.4	0.98		0.37	
				M	61			0.0	26.2	0.83		0.24	
PAKISTAN	Karachi National micronutrient survey	N.S.	0-14.99	F	N.S.	Z	?	16.0			3014		
				M			?	20.0					
		2-3/90	B	532	A	2.3	48.3			3027			
		1976-77	all	6738			4.5	13.0			3007		
				F			4.3	14.8					
				M			4.7	12.7					
			children	F	N.S.		0.0	2.1					
	pregnant	F	N.S.		5.6	10.7							
	B	154			?	57.0			3014				
	B	159			?	70.0							
	B	265			?	29.0							
SAUDI ARABIA	Riyadh	N.S.	N.S.	B	N.S.	Z	1.1	9.9			3009		
YEMEN	Tihuna Region - rural	6-7/92	1-5.99	B	319	Z	7.2	62.4			3025		
			1-1.99		60			5.0	65.0				
			2-2.99		58			13.8	62.1				
			3-3.99		71			11.3	66.2				
			4-4.99		56			3.6	60.7				
			5-5.99		74			2.7	58.1				
WESTERN PACIFIC REGION													
CHINA	National - Beijing (urban) Rural China	1982	0-2	B	N.S.	Z		17.0			6018		
		N.S.	5-2					20.0					
MALAYSIA	Peninsular Malaysia	1964	10-14	F	70	Z	5.7	22.8	0.97	0.05	6006		
				M	79			2.5	24.0	1.06		0.06	
			15-44	F	111			2.7	9.0	1.28		0.05	
				M	75			0.0	1.3	1.50		0.05	
			5-9	F	59			8.5	30.5	0.97		0.07	
				M	57			0.0	24.6	0.98		0.06	

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified
** Median

INDICATORS OF VITAMIN A DEFICIENCY - SERUM VITAMIN A											
Country	Geographic Area	Survey year	Age group	Sex	Sample size	*	Vitamin A Levels (µmol/l)				Ref. no.
							< 0.35	< 0.70	Mean	S.D.	
	Sungai Choh Ulu Rening Rural community - peninsular Malaysia	1977 1984	>45	F	24		4.2	25.0	1.21	0.13	
				M	22		0.0	4.5	1.41	0.12	
			children	F	25		4.0	16.0	1.02	0.07	
			pregnant		8		0.0	12.5	1.15	0.13	
			0-4	B	22			32.0	0.84	0.21	
			school		80			27.0	0.91	0.39	
			0-4		38			32.0	0.88	0.28	
			school		69			16.0	1.12	0.39	
			0-4		25			12.0	1.12	0.33	
			12-17.99	F	61			3.0	1.93	0.67	
				M	32			16.0	1.54	0.77	
			18-45	F	353			12.0	1.65	0.84	
				M	152			7.0	1.61	0.67	
			>46	F	14			7.0	1.47	0.60	
	M	14			0.0	1.89	1.16				
school	B	40			10.0	1.16	0.44				
MICRONESIA (FEDERATED STATES OF)	National - Chuuk	12/88-5/89	3-6.99	B	254	Z	15.0	64.0	0.60	6017	
PAPUA NEW GUINEA	East Sepik	N.S.	5-15	B	237	Z	50.0	91.0	0.41	0.20	6021
			5-5.99		74		2.7	58.1			
			5-6		N.S.			0.31	0.02		
			6-15					0.44	0.02		
PHILIPPINES	Cebu - 12 poor areas	1973-74	1-16	B	1700	Z	17.0	47.0			6012
	- pre-intervention non MSG group				727				0.68		
	- post-intervention non MSG group				727				0.55		
	- pre-intervention MSG group				387				0.74	0.43	
	- post-intervention MSG group				387				1.00	0.52	
	Cebu Island				1715		17.0	57.0			6002
	- rural-coastal				N.S.		20.0	62.0			
	- urban-squatter						17.0	54.0			
	- urban-barrio						15.0	59.0			
	Northern Mindanao	1986	5-99					20.0			7003
			1-6					15.7			
			children	F				10.0			
			pregnant					7.5			
	Southern Tagalog		0.5-99	B				19.5			
			1-5.99					10.4			
			children	F				1.4			
			pregnant					7.3			
	Western Visayas		0.5-99	B				33.3			
			1-5.99					10.9			
			children	F				21.0			
			pregnant					24.1			
	National	1987	0-5.99	B	3389		2.6				6022

* Laboratory Assessment: 'A' = HPLC, 'B' = Spectrophotometry, 'C' = Colorimetry, 'D' = Fluorescence, 'Z' = Not Specified
 ** Median

MDIS

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY

BIBLIOGRAPHIC REFERENCES

BIBLIOGRAPHIC REFERENCES

- 1002 Resnikoff S. Aspects épidémiologiques de la xérophtalmie au Tchad. *Médecine tropicale*, 1988, **48**: 27-32.
- 1004 De Sole G et al. Vitamin A deficiency in Southern Ethiopia. *American journal of clinical nutrition*, 1987, **45**: 780-784.
- 1006 *Assessment of Visual Status Among Ethiopian Refugees in Sudan*. Helen Keller International, January 1985.
- 1014 Foster A, Kavishe F, Sommer A, Taylor HR. A simple surveillance system for xerophthalmia and childhood corneal ulceration. *Bulletin of the World Health Organization*, 1986, **64**(5): 725-28.
- 1016 Madzima, M. Personal communication providing table with preliminary results of Vitamin A study in Zimbabwe. 7 August 1992.
- 1018 Ministry of Health. *Malawi Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 1021 *Lower Shire Valley Ocular Disease Survey-Final Report*. Malawi, Ministry of Health-Government of Malawi, 1986.
- 1022 Eastman SJ. *UNICEF Vitamin A Consultancy Report-Mali Country Review*. May 1986.
- 1023 Carlier C, Mouliat P, Cecon J F. Prevalence of malnutrition and vitamin A deficiency in the Diorbél, Fatick and Koalack regions of Senegal: A controlled study. *American journal of clinical nutrition*, 1991, **53**: 74-77.
- 1024 Kavishe F, Temalilwa C et al. *Towards a National Nutrition Blindness Programme in Tanzania. Report to the 8th IVACG Meeting*. Geneva, May 1984 (Project No. 2401 TFNC Report No. 867).
- 1025 van der Haar F. *Ten Year United Nations Action Programme - Prevention and Control of Vitamin A Deficiency in Tanzania - Tentative FAO Elements*. 1986.
- 1027 Pepping F, van der Giezen A. Food Consumption of Children with and without Xerophthalmia in Rural Tanzania. *Tropical and Geographical Medicine*, 1989, **41**:14-21.
- 1028 Eastman SJ. *UNICEF Vitamin A Consultancy Report - Tanzania Country Review*. July 1986.
- 1029 Sheffield VM. *Report on Vitamin A Supplementation Programs - Ethiopia*. New York, Helen Keller International, 1985.
- 1030 Bonnet S, Resnikoff S, Casten R et al. *Assessment of Vitamin A Status in a Rural Area in Mali*. Unpublished draft, 1989.

- 1031 McLaren DS ed. VAST (Vitamin A Supplementation Trials) in Ghana. *Xerophthalmia Club - Bulletin* 4B, October 1991.
- 1032 Kawuma M, Sserunjogi L. *Kamuli Blindness and Vitamin A Deficiency Survey*. 1992 (Ministry of Health Technical Report, Series 1, No.1).
- 1033 *The Annual Report for the Malawi Vitamin A Project*. Save the Children. 1988-89.
- 1034 Sukwa T, Mwandu D, Kapui A et al. The Prevalence and Distribution of Xerophthalmia in Preschool Children of the Luapala Valley, Zambia. *Journal of tropical pediatrics*, 1988, **34**: 12-15.
- 1035 *Enfants et Femmes au Burkina Faso - Une Analyse de la Situation*. Ouagadougou, UNICEF, Septembre 1991.
- 1037 *Iodine Deficiency Disorders in Namibia and Data on the Status of Vitamin A and Iron. Draft Report*. Windhoek, Ministry of Health and Social Services, Government of the Republic of Namibia, December 1992.
- 1038 Ministry of Health. *Cape Verde Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 1039 Steinkuller PG. Nutritional Blindness in Africa. *Social science and medicine*, 1983, **17**(22): 1715-1721.
- 1040 Ministry of Health. *Guinea Country Report*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 1041 Ministry of Health. *Benin Country Report*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 1042 Ministry of Health. *Mauritania Country Report*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 1043 Ette SI, Esenowe EJ. Plasma concentrations of retinol and vitamins E and C during the first twelve months of post natal life. *Tropical and geographical medicine*, 1986, **38**: 1-5.
- 1044 Coutsooudis A, Mametjas D. Vitamin A Deficiency among children in a periurban South African settlement. *American journal of clinical nutrition*, 1993, **57**:904-907.
- 1049 Atina E, Wilson R et al. *Prevalence of Xerophthalmia and Risk of Vitamin A Deficiency among children in the extreme north of Cameroon*. 1992.
- 1050 Nathanail L, Powers H. Vitamin A Status of Young Gambian Children: biochemical evaluation and CIC. *Annals of tropical paediatrics*, 1992, **12**:67-73.

- 1051 Morris Saul. Personal correspondence providing data from the Ghana VAST study.
- 1052 Zewdie Wolde-Gabriel. *Micronutrient deficiencies in Ethiopia and their inter-relationships*. Den Haag, CIP-Data Koninklijke Bibliotheek, 1992.
- 1053 Carlier C et al. Assessment of the vitamin A status of preschool and school age Senegalese children during a cross-sectional study. *International journal of vitamin and nutritional research*, 1991, 62:209-215.
- 1055 *National Survey on Iodine, Vitamin A and Iron Status of Women and Children in Lesotho*. Maseru, Food and Nutrition Coordinating Office, Ministry of Health/UNICEF Maseru, 1994.
- 1056 Sturchler D, Holzer B, Hanck A, Degremont A. The influence of schistosomiasis on the serum concentrations of retinol and retinol binding protein of a rural population in Liberia. *Acta tropica* 1983, 40:261-269.
- 1057 *Chad Country Report. Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 1058 Arthur P. *Prevalence of Vitamin A Deficiency in Kintampo District: Report of a Survey*. London, London School of Hygiene and Tropical Medicine, 1994.
- 1059 Roy C. *Parametres Biologiques des Enfants Malnutris (MPE) et d'un Groupe d'Enfants à Status Nutritionnel Correct Identifiés Lors de l'Enquête Nutritionnelle Transversale Réalisée dans le Secteur de Dabakala en Côte d'Ivoire*. Abidjan, Institut National de Santé Publique, Mai 1990.
- 1061 *Magnitude and Distribution of Vitamin A Deficiency in Nigeria*. VITAL. Data presented by NMH Thailand, 1994.
- 1063 Tebi A et al. *Evaluation du Statut en Vitamine A de l'Enfant de 6 à 59 Mois du Nord Ouest de la Côte d'Ivoire*. Abidjan, Institut National de Santé Publique, 1994.
- 1064 Ghana VAST Study Team. Vitamin A supplementation in northern Ghana: effects on clinic attendances, hospital admissions, and child mortality. *Lancet*, 1993, 342: 7-12.
- 1065 Wolde-Gabriel Z et al. Interrelationship between vitamin A, iodine and iron status in schoolchildren in Shoa Region, Central Ethiopia. *British journal of nutrition*, 1993, 70: 593-607.
- 1067 Lechat MF et al. Epidémiologie de l'avitaminose A au Niger. *Annals de société belge de médecine tropicale*, 1976, 56(4-5): 333-342.
- 1068 Gutekunst R. Tables 10-12 from an unknown source. Quotes data for Namibia.

- 1069 Borel E and Etard J. Carences en vitamine A dans une population rurale de Mauritanie et absence de corrélation avec la schistosomiase urinaire. In: *Troisième Journées Scientifiques Internationales du GERM*. Nianing, 1987, 356-363.
- 1070 Samba C, Galan P, Luzeau R, Amedee-Manesme O. Vitamin A deficiency in pre-school age Congolese children during malarial attacks. *International journal of vitamin and nutritional research*, 1990, **60**: 215-223.
- 1071 Carlier C et al. Efficacy of massive oral doses of retinyl palmitate and mango (*Mangifera indica* L.) consumption to correct an existing vitamin A deficiency in Senegalese children. *British journal of nutrition*, 1992, **68**: 529-540.
- 1072 Wolde-Gebriel Z et al. Severe vitamin A deficiency in a rural village in the Hararge region of Ethiopia. *European journal of clinical nutrition*, 1993, **47**: 104-114.
- 1073 Carlier C et al. Annual assessment of the vitamin A and nutritional status of children during two cross-sectional surveys. *International journal of vitamin and nutritional research*, 1992, **62**: 216-220.
- 1074 Rankins J et al. Undernutrition and vitamin A deficiency in the Department of Linguère, Louga Region of Senegal. *American journal of clinical nutrition*, 1993, **58**: 91-97.
- 1075 Julien M et al. *The Assessment of Vitamin A Deficiency in Three Cities in Mozambique*. Paper presented at the XV IVACG Meeting, Arusha, 1993.
- 2002 Pancheco Santos LM et al. *A Preventive Program for Vitamin A Deficiency in Paraiba, Northeast Brazil*. Paper presented at the Xth IVACG Meeting, Hyderabad, 1985.
- 2004 *National Committee for the Blind and Deaf of Guatemala/UNICEF. Report on the Joint Ophthalmic and Nutritional Survey and Delivery of Services in Nebaj, Quiche 16-25 May 1984*. 1985.
- 2007 Ministério da Saúde-Instituto Nacional de Alimentação e Nutrição - INAN. *Hypovitaminosis A in Brazil*. Paper presented to IVACG, Geneva, May 1977.
- 2008 Favoro RMD et al. Vitamin A Status of Young Children in Southern Brazil. *American journal of clinical nutrition*, 1986, **43**: 852-858.
- 2009 Santos LM et al. Xerophthalmia in the state of Paraiba, northeast Brazil: clinical findings. *American journal of clinical nutrition*, 1983, **38**: 139-144.
- 2010 Arroyave G, Mejia LA, Aguilar JK. The effect of vitamin A fortification of sugar on the serum vitamin A levels of preschool Guatemalan children: A longitudinal evaluation. *American journal of clinical nutrition*, 1981, **34**: 41-49.

- 2011 Pilch SM. Analysis of Vitamin A Data from the Health and Nutrition Examination Surveys. *Journal of Nutrition*, 1987, **117**: 636-640.
- 2014 Center for Disease Control. Unpublished Data from NHANES I, 1971-74.
- 2015 *Estrategias de Intervención para Hipovitaminosis*. San Salvador, Calma Centro de Arroyo de Lactancia Materna, 1990.
- 2016 *Encuesta de Vitamina A y Encuesta de Consumo - Áreas Deprimidas*. Bolivia, VITAL/USAID, 1991.
- 2021 Makdani D, Rizner J, et al. Vitamin A and Zinc Nutriture of Children in Belize, Central America. *FASCB journal*, 1991, A953.
- 2024 *Brazil Country Assessment. Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 2025 *Honduras Country Assessment. Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 2026 *Deficiencia de Vitamina A en Provincias de Pobreza Crítica del Ecuador*. Quito, Ministerio de Salud Pública del Ecuador, Febrero 1994.
- 2027 *Vitamin A Field Support Project - VITAL. Trip Report Panama December 7-11, 1993*. Washington, D.C., 1993 (USAID Report No. TR-21).
- 2028 de Caballero E. *The Prevalence of Vitamin A Deficiency and Iron Deficiency Anemia of Preschool Children in Panama*. Panama, Ministry of Public Health.
- 2029 Dricot d'Ans C et al. Geographic distribution of xerophthalmia in the state of Paraíba, Northeast Brazil. *Ecololgy of Food and Nutrition*, 1988, **22**: 131-138.
- 2031 *Deficiencia de Vitamina A en la Región del Suroeste de la República Dominicana*. Santa Domingo, Centro Nacional de Investigaciones en Salud Materno Infantil (CENISMI), 1991.
- 2032 Santos LMP et al. *Avaliacao Antropometrica, Hipovitaminose A e Anemia em pre-Escolares do Semi-Arido da Bahia*. Written in conjunction with USAID, project no. 598-06-G-SS-7003-00.
- 2033 *Encuesta Nacional de Vitamin A*. República de Panamá, Ministerio de Salud, Departamento de Nutrición y Dietética, 1992.
- 2034 Salazar-Lindo E, Salazar M and Alvarez JO. Association of diarrhea and low serum retinol in Peruvian children. *American journal of clinical nutrition*, 1993, **58**: 110-113.

- 3003 Ibrahim K et al. Plasma vitamin 'A' and carotene levels in Karachi population. *Journal of Pakistani medical association*, 1987, 37:117-126.
- 3005 Ministry of Health. *Oman Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3006 Ministry of Health. *Pakistan Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3007 Eastman SJ. *UNICEF Vitamin A Consultancy - Pakistan Country Report*. May 1986.
- 3008 Ministry of Health. *Morocco Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3009 Alshoshan AA. *Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3010 Ministry of Health. *Sudan Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3011 Sheffield V. *Report on Vitamin A Supplementation Programs*. New York, Helen Keller International, 1986.
- 3012 Ministry of Health. *Yemen Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 3013 Resnikoff S et al. Assessment of Vitamin A deficiency in the Republic of Djibouti. *European journal of clinical nutrition*, 1991.
- 3014 UNICEF. *Vitamin A Deficiency in Pakistan - Report of the Status of Knowledge on Prevalence of Vitamin A Deficiency in Pakistan*. Pakistan, Aga Khan University Hospital, King Edward Medical College, December 1990.
- 3016 *Situation Analysis of Women and Children in the Islamic Republic of Iran*. Teheran, UNICEF, 1992.
- 3017 Pigot D. *Health and Nutrition in Kabul, Herat, and Mazar-I-Sharif - A Preliminary Assessment*. September 1989.
- 3019 Abbas KA, Reddy SK. *Assignment Report: Appraisal of Vitamin A Deficiency - Sultanate of Oman*. 1981.
- 3020 Rosen DS et al. *Xerophthalmia in the Tihama Region of Yemen, Preliminary Analysis of a Prevalence Survey*. 1992.

- 3021 Bloem M, Farooq S, Kultab A. *Vitamin A Deficiency and Malnutrition in Southern Iraq - Rapid Assessment Report*. HKI/SC/UNICEF, May 1991.
- 3022 *Vitamin A Deficiency and Xerophthalmia Prevalence Assessment Survey in Northern Darfur*. Helen Keller International, February-March 1988:
- 3024 *Vitamin A Survey: Red Sea Province*. New York, Helen Keller International, 1989.
- 3025 Pizzarello L. Table 3 Serum Retinol Levels by Age in Rural Tihana. Personal communication from S Burger (HKI), 10 June 1993.
- 3026 Alnwick D. *Report of a Review of UNICEF Assistance to Achieve Nutrition Mid-Decade Goals in Iraq*. Baghdad, UNICEF, November 1994.
- 3027 Molla A et al. Vitamin A status of children in the urban slums of Karachi, Pakistan, assessed by clinical, dietary, and biochemical methods. *American journal of tropical medical hygiene*, 1993, **48**(1): 89-96.
- 3028 Resnikoff S. Personal correspondence providing Table 4 Plasma retinol levels among children 4 to 6 years old from Djibouti. 1988.
- 3029 *National Study on the Prevalence of Vitamin A Deficiency (VAD) Among Children 6 Months to 7 Years*. Sultanate of Oman Ministry of Health, WHO/UNICEF, 1995.
- 4001 Glick Z, Reshef A. Vitamin A Status and related nutritional parameters of children in East Jerusalem. *American journal of clinical nutrition*, 1973, **26**: 1229-1233.
- 4002 Wetherilt H et al. Blood vitamin and mineral levels in 7-17-year-old Turkish children. *International journal of vitamin and nutritional research*, 1992, **62**: 21-29.
- 4003 *Romania Country Report. Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 4004 Hatun S, Tezic T. *Vitamin A Status of Healthy Infants in Ankara, Turkey*. Ankara, Dr. Sami Ulus Children's Hospital, 1994.
- 4005 Malvy JMD et al. Retinol, *B*-carotene and *a*-tocopherol status in a French population of healthy children. *International journal of vitamin and nutrition research*, 1989, **59**: 29-34.
- 4006 Morse C. *A Report on the Prevalence of Anemia in Muynak District, Karakalpakstan, Uzbekistan*. Crosslink International, September 1993.
- 5002 *Bangladesh Nutritional Blindness Survey, 1982-83*. Dhaka, Helen Keller International, 1985.

- 5003 Working Group on IDD Control, Myanmar. *Available Data on Vitamin A Status of Myanmar Children*. Paper presented at Interdepartmental Workshop on Prevention and Control of Vitamin A Deficiency in Myanmar, 29 March 1990.
- 5007 Indirabai K, Bhatt JV, Vaidyanathan K. Epidemiological survey for prevalence of xerophthalmia. *Indian pediatrics*, 1986, **23**: 135-139.
- 5012 Xerophthalmia surveillance - prevalence of xerophthalmia in Lombok. *WHO Weekly Epidemiological Record*, 1984, **17**: 129-130.
- 5016 Muhilal et al. Vitamin A-fortified monosodium glutamate and health, growth, and survival of children: A controlled field trial. *American journal of clinical nutrition*, 1988, **48**: 1271-1276.
- 5017 Tarwotjo I et al. Dietary practices and xerophthalmia among Indonesian children. *American journal of clinical nutrition*, 1982, **35**: 574-581.
- 5021 Djunaedi E et al. Impact of vitamin A supplementation on xerophthalmia: A randomized controlled community trial. *Archives of ophthalmology*, 1988, **106**: 218-222.
- 5026 Atukorala S. Control of Vitamin A Deficiency and Studies on Vitamin A in Sri Lanka. In: *Proceedings-Regional Meeting in Vitamin A, Jakarta, November 3-5, 1988*. WHO/MOH (Indonesia), 1988.
- 5027 Brink EW et al. Vitamin A status of children in Sri Lanka. *American journal of clinical nutrition*, 1982, **35**: 574-581.
- 5029 Bloem MW et al. A prevalence study of Vitamin A deficiency and xerophthalmia in Northeastern Thailand. *American journal of epidemiology*, 1989, **129**(6): 1095-1103.
- 5030 Bloem M.W. *Vitamin A Deficiency, Anemia and Infectious Diseases in Northeast Thailand* [Thesis]. Utrecht, November 1988.
- 5032 Khan MU, Haque E, Khan MR. Nutritional ocular diseases and their association with diarrhea in Matlab, Bangladesh. *British journal of nutrition*, 1984, **52**: 1-9.
- 5033 Ministry of Health. *Bhutan Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 5034 Tilden R. *Prevention and Control of Vitamin A Deficiency, Xerophthalmia and Nutritional Blindness in Bhutan - Assignment Report, 3-12 March 1986*. New Delhi, 1986 (WHO Project ICP NUT 005).
- 5035 Rahmathullah L et al. Reduced mortality among children in Southern India receiving a small weekly dose of Vitamin A. *New England journal of medicine*, 1990, **323**: 929-935.

- 5036 Sharma SK et al. Xerophthalmia in preschool children. *Indian pediatrics*, 1987, **24**: 645-650.
- 5037 Vijayaraghavan K et al. Effect of massive dose of Vitamin A on morbidity and mortality in Indian children. *Lancet*, 1990, **336**: 1342-1345.
- 5040 Garg S et al. Vitamin A Deficiency in pre-school children. *Indian pediatrics*, 1984, **21**: 491-494.
- 5041 Shah P.M. *Strategies for Prevention of Malnutritional Blindness in India: Operational Methodology, Management, Monitoring and Cost Benefit*. A Report Prepared for the Commonwealth Society for the Blind, July 1978.
- 5042 Eastman SJ. *India Country Review - Vitamin A*. UNICEF Vitamin A Consultancy - April 23, 1986.
- 5043 *India Country Paper. Quotes NWMB Data*. International Conference on Nutrition. Rome, Ministry of Health/FAO/WHO, December 1992.
- 5044 Pant I, Gopaldes TA. *Studies on Vitamin A Deficiency in Underprivileged Children (5-15 years)*. Baroda, University Department of Food and Nutrition.
- 5045 Gopaldes TA. *Nutrition Health Education Program to Combat Vitamin A Deficiency in Gujarat*. Baroda, University Department of Food and Nutrition.
- 5046 Kusin JA, Sinaga HS, Marpaung AM. Xerophthalmia in North Sumatra. *Tropical and geographical medicine*, 1977, **29**: 41-46.
- 5047 Ministry of Health. *Indonesia Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 5048 Xerophthalmia Surveillance. *WHO weekly epidemiology record*, 1994, **17**: 129-130.
- 5050 Kusin JA et al. Anemia and hypovitaminosis A among rural women in East Java, Indonesia. *Tropical and geographical medicine*, 1980, **32**: 30-39.
- 5051 Ministry of Health. *Myanmar Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 5052 *Children and Women in Nepal: A Situation Analysis 1992*. Kathmandu, National Planning Commission, HMG, UNICEF Nepal, 1992.
- 5054 Brink EW et al. Vitamin A Status of Children in Sri Lanka. *American journal of clinical nutrition*, 1979, **32**: 84-91.
- 5055 Ministry of Health. *Sri Lanka Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.

- 5057 Cohen N et al. Impact of massive doses of vitamin A on nutritional blindness in Bangladesh. *American journal of clinical nutrition*, 1987, **45**:970-976.
- 5058 Devadas R. *Dietary Supplementation and Assessment of Vitamin A Nutrition*. Unpublished draft.
- 5059 Pandey MR et al. Reduction in total under-five mortality in Western Nepal through community based antimicrobial treatment of pneumonia. *Lancet*, 1991, **338**: 993-997.
- 5060 Bergen HR et al. Vitamin A and vitamin E status of rural preschool children in West Java Indonesia and their response to oral doses of vitamin A and of vitamin E. *American journal of clinical nutrition*, 1988, **48**: 279-285.
- 5061 West KP. *Report of a Consultancy on Vitamin A Deficiency in Myanmar*. UNICEF and Government of Myanmar, May 1990.
- 5062 Tilden R. *Prevention and Control of Vitamin A Deficiency, Xerophthalmia and Nutritional Blindness in Sri Lanka*, New Delhi, WHO, 1986 (Assignment report 2/15/86-3/1/86).
- 5063 Thein D, Myint T. *Vitamin A Status of children in Myanmar*. Paper presented at Interdepartmental Workshop on Prevention and Control of Vitamin A Deficiency. Myanmar, National Nutrition Center.
- 5064 Shankar M. *Vitamin A Activity in USAID - Assisted Integrated Child Development Services program in Panchmahals district, Gujarat, and Chandrapur district, Maharashtra*. August 1991.
- 5065 *The Epidemiology of Blindness in Nepal*. Michigan, Seva Foundation, 1988.
- 5068 Mathews T. *Vitamin A Deficiency and its Prevention in Surkhet District*. Unpublished draft paper.
- 5070 *Report on the Prevalence of Inadequate Vitamin A Nutriture in Preschool Children of North and Northeast Thailand*. Bangkok, Mahidol University, 1991.
- 5071 Kolhede C. Unpublished data available to WHO from MORVITA Trials, 1992.
- 5072 Varavithaya W et al. Vitamin A status of children with measles. *Tropical and geographical medicine*, 1986, **38**: 359-361.
- 5073 Bloem MW. *Vitamin A deficiency, anemia and infectious disease in Northeast Thailand* [Unpublished Thesis draft]. 1988.
- 5074 *An Evaluation of the Vitamin A Deficiency Prevention Pilot Project in Indonesia 1973-75*. American Foundation for Overseas Blind, Inc.
- 5075 Udomkesmalee E et al. Effect of Vitamin A and zinc supplementation on the nutriture of children in northeast Thailand. *American journal of clinical nutrition*, 1992, **56**: 50-57.

- 5078 Anderson J, Tilden R. *Prevention and Control of Vitamin A Deficiency, Xerophthalmia, and Nutritional Blindness in Bangladesh*. Report of a Joint FAO/WHO mission to Bangladesh, 18 March - 12 April 1986.
- 5079 Udomkesmalee E. Unpublished information presenting biochemical data for Thailand, 1991.
- 5080 *Report of Xerophthalmia Prevalence Surveys in Five Districts of the Far and Mid-Western Regions of Nepal*. Vitamin A Field Support Project - VITAL, June-September 1993.
- 5081 Udomkesmalee E et al. *Assessment of Marginal Vitamin A Deficiency in Thai School Children by Impression Cytology, Dark Adaptometry and Serum Retinol*. Bangkok, Institute of Nutrition, Mahidol University.
- 5082 Valyasevi A, Charoenkiatkul S. *Dietary Approaches for Prevention of Vitamin A Deficiency*. A paper prepared for the 8th IVACG Meeting in Geneva, Switzerland, June 19-23, 1984. Bangkok, Institute of Nutrition, Mahidol University.
- 5083 Tanumihardjo S et al. Assessment of the vitamin A status in lactating and nonlactating, nonpregnant Indonesian women by use of the modified-relative-dose-response (MRDR) test. *American journal of clinical nutrition*, 1994, **60**: 142-147.
- 5084 Hussain A et al. Determinants of night blindness in Bangladesh. *International journal of epidemiology*, 1993, **22**(6) 1119-1125.
- 5085 Gopaldas T, Gujral S, Abbi R. Prevalence of xerophthalmia and efficacy of vitamin A prophylaxis in preventing xerophthalmia co-existing with malnutrition in rural Indian children. *Journal of tropical pediatrics*, 1993, **39**: 205-208.
- 5086 Henning A et al. Vitamin A deficiency and corneal ulceration in south-east Nepal: implications for preventing blindness in children. *Bulletin of the World Health Organization*, 1991, **69**(2): 235-239.
- 6002 Solon FS et al. Vitamin A deficiency in the Philippines: a study of xerophthalmia in Cebu. *American journal of clinical nutrition*, 1978, **31**: 360-368.
- 6006 Ministry of Health. *Malaysia Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 6010 van der Haar F, ten Doesschate J. *Report on an advisory mission to Viet Nam 22nd May to 8 June 1984 on the subject of Vitamin A Deficiency and Xerophthalmia*. Subproject of Nutrition (1983-85) Inter-University Medical Cooperation of the Netherlands and Vietnam, Wageningen, July 1984.
- 6011 Dricot J. *The Vitamin A Deficiency Control Programme in the Socialist Republic of Viet Nam*. Hanoi, UNICEF, May 1988.

- 6012 Solon FS et al. Planning, implementation and evaluation of a fortification program. Control of Vitamin A deficiency in the Phillipines. *Journal of the american dietetic association*, 1979, 74: 112-118.
- 6013 *Kiribati Vitamin A Deficiency Assessment*. The Foundation for the Peoples of the South Pacific, September-October 1989.
- 6014 *Solomon Islands Xerophthalmia Survey*. Vitamin A Field Support Project - VITAL, 1992 (Report No. TA-8).
- 6015 Giay T, Khoi HH, Nhan NT. Vitamin A Deficiency in Preschool Age Children in Vietnam Today. *Xerophthalmia*.
- 6016 Ministry of Health. *Vietnam Country Paper*. FAO/WHO International Conference on Nutrition, Rome, December 1992.
- 6017 Lloyd-Puryear M et al. Vitamin A deficiency in Micronesia: A statewide survey in Chuuk. *Nutritional Research*, 1991, 11:1101-1110.
- 6018 *China Country Paper. Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 6019 Danks J, Kaufman D, Rait J. A clinical and cytological study of vitamin A deficiency in Kiribati. *Australian and New Zealand journal of ophthalmology*, 1992, 20(3): 215-218.
- 6020 Feldon K et al. *Preliminary Results of HKI/MOH Vitamin A Deficiency Surveys*. May-August 1993.
- 6021 Genton B et al. *Vitamin A Status and Malaria Infection, Morbidity, and Immunity in a Highly Endemic Area of Papua New Guinea*. XVI IVACG Meeting Abstract Form.
- 6022 *Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991. Presents data for Philippines*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 6023 Ophthalmic Morbidity Survey in Kiritimati. Personal correspondence from CLO Kiribati. 5 August 1994.
- 6024 Velandria F et al. *Fourth National Nutrition Survey*. Philippines, Food and Nutrition Research Institute, 1993.
- 6025 *Tuvalu Xerophthalmia Survey*. Vitamin A Field Support Project - VITAL, June 4-27, 1991 (Report No. TA-7).

- 6026 *National Vitamin A Survey - Lao PDR, 1995. Final Report.* Vientiane, Ministry of Health/World Health Organization/UNICEF, 1995.
- 6027 Bloem M, Gorstein J. *Viet Nam: Xerophthalmia Free -1994 National Vitamin A Deficiency and Protein-Energy Malnutrition Prevalence Survey.* 5-17 March, 1995.
- 6029 Greenop P, Dignan C. *1994 Report of the National Nutrition Survey of the Republic of the Marshall Islands.* August, 1994.
- 7001 *Vitamin A Facts - African region (Selected Countries).* Vitamin A Field Support Project - VITAL. Washington, D.C., prepared for the Office of Nutrition - Bureau of Science and Nutrition, USAID, 1992 (Report No. IN-7).
- 7002 *Vitamin A Facts - Latin America and Caribbean Region (Selected Countries).* Vitamin A Field Support Project - VITAL. Washington, D.C., paper prepared for the Office of Nutrition, USAID, 1992 (Report No. IN-7).
- 7003 *Vitamin A Facts-Asia Region (Selected Countries).* Vitamin A Field Support Project-VITAL. Washington D.C., prepared for the Office of Nutrition-Bureau of Science and Nutrition-USAID, 1992 (Report No. IN-7).
- 7004 Arthur P. Vitamin A Deficiency in Africa. *Vital News*, 1992, 3(2):1-4.
- 7005 Udomkesmalee E. Vitamin A Deficiency in Asia. *Vital News*, 1992, 3(3):1-7.
- 7007 Pan American Health Association-Food and Nutrition Program. Summary Data compiled by WHO/PAHO.
- 7008 *Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991. Presents data for Indonesia, Bhutan and Ghana.* Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 7009 Plan of Action for the Elimination of Vitamin A Deficiency from the Americas. Provisional Agenda Item 4.3. PAHO/WHO, June 1992.
- 7010 Mora J. *Simposio Sobre la Deficiencia de Vitamina A en América Latina y el Caribe: Situación Actual y Perspectiva de Solucion. Situación Actual de la Deficiencia de Vitamina A en América Latina y el Caribe.*
- 7011 Eastman SJ. UNICEF Vitamin A Consultancy - *Central America Vitamin A Review - DRAFT.* 18 August 1986.
- 7012 Le Francois P et al. Vitamin A Status in Three West African Countries. *International journal of vitamin and nutritional research*, 1980, 50(4): 352-363.

- 7015 *Xerophthalmia and Trachoma in Burkina Faso, Chad, Mali and Niger*. Unpublished report. Helen Keller International Inc., 1986.
- 7016 *Databank on Vitamin A Deficiency Disorders for the WHO African Region*. See individual country papers for specific references. WHO Regional Office for Africa, May 1992.
- 7018 *Conference on Ending Hidden Hunger. Proceeding of a policy conference on micronutrient malnutrition, Montreal, October 1991. Presents data for Angola*. Atlanta, GA, Task Force for Child Survival and Development, 1991.
- 7020 *Proceedings of an Inter-Agency Coordination Meeting on Micronutrient Deficiency Control in Washington, D.C.* WHO/PAHO, April, 1993.
- 7021 *Tercer Taller Regional Sobre Deficiencias de Vitamina A y Otros Micronutrientes en América Latina y el Caribe, Recife (Brasil), agosto 23-27, 1993. Situación de Vitamina A, Hierro y Yodo en la Región*. Virginia, Vitamin A Field Support Project - VITAL, November 1993.
- 7022 *Towards Comprehensive Programs to Reduce Vitamin A Deficiency*. Report of the XV International Vitamin A Consultative Group Meeting. Tanzania, 1993.
- 7023 *Review of USAID/VITAL - Supported Vitamin A Deficiency Surveys*. Vitamin A Field Support Project - VITAL. Prepared for the Office of Nutrition - Bureau of Science and Nutrition - USAID, Washington D.C., 1994 (Report No. FA-24).
- 7024 Miller D. Personal communication received 17 February 1993.

MDIS

GLOBAL PREVALENCE OF VITAMIN A DEFICIENCY

ANNEXES

LISTING OF COUNTRIES ACCORDING TO WHO AND UNICEF REGIONS

WHO REGIONS

Africa	Americas	South-East Asia	Europe	Eastern Mediterranean	Western Pacific
Algeria	Antigua & Barbuda	Bangladesh	Albania	Afghanistan	Australia
Angola	Argentina	Bhutan	Armenia	Bahrain	Brunei Darussalam
Benin	Bahamas	Democratic People's Republic of Korea	Austria	Cyprus	Cambodia
Botswana	Barbados	India	Azerbaijan	Djibouti	China
Burkina Faso	Belize	Indonesia	Belarus	Egypt	Cook Islands
Burundi	Bolivia	Maldives	Belgium	Iran (Islamic Rep. of)	Fiji
Cameroon	Brazil	Mongolia	Bosnia & Herzegovina	Iraq	Japan
Cape Verde	Canada	Myanmar	Bulgaria	Jordan	Kiribati
Central African Republic	Chile	Nepal	Croatia	Kuwait	Lao P.D.R.
Chad	Colombia	Sri Lanka	Czech Republic	Lebanon	Malaysia
Comoros	Costa Rica	Thailand	Denmark	Libyan Arab Jamahiriya	Marshall Islands
Congo	Cuba		Estonia	Morocco	Micronesia (Fed. States of)
Côte d'Ivoire	Dominica		Finland	Oman	Nauru
Equatorial Guinea	Dominican Repub.		France	Pakistan	New Zealand
Eritrea	Ecuador		Georgia	Qatar	Niue
Ethiopia	El Salvador		Germany	Saudi Arabia	Papua New Guinea
Gabon	Grenada		Greece	Somalia	Philippines
Gambia	Guatemala		Hungary	Sudan	Republic of Korea
Ghana	Guyana		Iceland	Syrian Arab Rep.	Samoa
Guinea	Haiti		Ireland	Tunisia	Singapore
Guinea-Bissau	Honduras		Israel	United Arab Emirates	Solomon Islands
Kenya	Jamaica		Italy	Yemen	Tokelau
Lesotho	Mexico		Kazakhstan		Tonga
Liberia	Nicaragua		Kyrgyzstan		Tuvalu
Madagascar	Panama		Latvia		Vanuatu
Malawi	Paraguay		Lithuania		Viet Nam
Mali	Peru		Luxembourg		
Mauritania	Puerto Rico		Malta		
Mauritius	St. Kitts & Nevis		Monaco		
Mozambique	St. Lucia		Netherlands		
Namibia	St. Vincent & the Grenadines		Norway		
Niger	Suriname		Poland		
Nigeria	Trinidad & Tobago		Portugal		
Rwanda	United States of America		Republic of Moldova		
Sao Tome & Principe	Uruguay		Romania		
Senegal	Venezuela		Russian Federation		
Seychelles			San Marino		
Sierra Leone			Slovakia		
South Africa			Slovenia		
Swaziland			Spain		
Togo			Sweden		
Uganda			Switzerland		
United Republic of Tanzania			Tajikistan		
Zaire			The former Yugoslav Rep. of Macedonia		
Zambia			Turkey		
Zimbabwe			Turkmenistan		
			Ukraine		
			U.K. of Great Britain & Northern Ireland		
			Uzbekistan		
			Yugoslavia		

UNICEF REGIONS

Eastern and Southern Africa	Central and West Africa	Middle East and North Africa	East Asia and the Pacific	Americas and the Caribbean	Developed/Industrialized Countries
Angola	Benin	Algeria	Brunei Darussalam	Antigua & Barbuda	Albania
Botswana	Burkina Faso	Bahrain	Cambodia	Argentina	Armenia
Burundi	Cameroon	Cyprus	China	Bahamas	Australia
Comores	Cape Verde	Djibouti	Cook Islands	Barbados	Austria
Ethiopia	Central African Rep.	Egypt	Fiji	Belize	Azerbaijan
Kenya	Chad	Iran, Islamic Rep.	Indonesia	Bolivia	Belarus
Lesotho	Congo	Iraq	Kiribati	Brazil	Belgium
Madagascar	Côte D'Ivoire	Jordan	Korea, Dem.	Chile	Bosnia/Herzegovina
Malawi	Equatorial Guinea	Kuwait	Korea, Rep. of	Colombia	Bulgaria
Mauritius	Gabon	Lebanon	Lao P.D.R.	Costa Rica	Canada
Mozambique	Gambia	Libya	Malaysia	Cuba	Croatia
Namibia	Ghana	Morocco	Marshall Islands	Dominica	Czech Republic
Rwanda	Guinea	Oman	Micronesia	Dominican Republic	Denmark
Seychelles	Guinea-Bissau	Qatar	Mongolia	Écuador	Estonia
Somalia	Liberia	Saudi Arabia	Myanmar	El Salvador	Finland
South Africa	Mali	Sudan	Papua New Guinea	Granada	France
Swaziland	Mauritania	Syria	Philippines	Guatemala	Georgia
Tanzania, UR	Niger	Tunisia	Samoa	Guyana	Germany
Uganda	Nigeria	Turkey	Singapore	Haiti	Greece
Zambia	Sao Tome & Principe	United Arab Emirates	Solomon Islands	Honduras	Hungary
Zimbabwe	Senegal	Yemen	Thailand	Jamaica	Iceland
	Sierra Leone		Tonga	Mexico	Israel
	Togo		Vanuatu	Nicaragua	Italy
	Zaire		Viet Nam	Panama	Japan
				Paraguay	Kazakhstan
				Peru	Kyrgyzstan
				St. Christ. & Nevis	Latvia
				St. Lucia	Lithuania
				St. Vincent & Gren.	Luxembourg
				Suriname	Malta
				Trinidad & Tobago	Moldova
				Uruguay	Netherlands
				Venezuela	New Zealand
					Norway
					Poland
					Portugal
					Romania
					Russian Federation
					San Marino
					Slovak
					Slovenia
					Spain
					Sweden
					Switzerland
					Tajikistan
					Turkmenistan
					Ukraine
					United Kingdom
					USA
					Uzbekistan
					Yugoslavia, Former

SELECTED WHO PUBLICATIONS AND DOCUMENTATION OF RELATED INTEREST

MICRONUTRIENT MALNUTRITION

Sommer A. *Vitamin A deficiency and its consequences: a field guide to their detection and control*. Third edition. Geneva, World Health Organization, 1995. Sw.fr. 17.- (11.90).

Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. Document NUT/94.7.

Global Prevalence of Iodine Deficiency Disorders. Micronutrient Deficiency Information System, Working Paper No.1. Geneva, World Health Organization, 1993.

INFANT AND YOUNG CHILD NUTRITION

Protecting, promoting and supporting breast-feeding: the special role of maternity services. A Joint WHO/UNICEF Statement. Geneva, World Health Organization, 1989. Sw.fr. 6.- (4.20).

Breast-feeding: the technical basis and recommendations for action. Document WHO/NUT/MCH/93.1.

Infant feeding: the physiological basis. Akre J. (ed.) Supplement to Vol. 67 (1989) of the *Bulletin of the World Health Organization*. Geneva, World Health Organization, 1990. Sw.fr. 20.- (14.-).

International Code of Marketing of Breast-milk Substitutes. Geneva, World Health Organization, 1981. Sw.fr. 4.- (2.80).

NUTRITIONAL ASSESSMENT, MONITORING

Measuring change in nutritional status. Guidelines for assessing the nutritional impact of supplementary feeding programmes for vulnerable groups. Geneva, World Health Organization, 1983. Sw.fr. 14.- (9.80).

An evaluation of infant growth. Summary of analyses performed in preparation for the WHO Expert Committee on physical status and the use and interpretation of anthropometry. Document WHO/NUT/94.8.

de Onis M et al. The worldwide magnitude of protein-energy malnutrition: an overview from the WHO Global Database on Child Growth. *Bulletin of the World Health Organization*, 1995, 71(6): 703-712.

WHO Working Group on Infant Growth. An evaluation of infant growth: the use and interpretation of anthropometry in infants. *Bulletin of the World Health Organization*, 1995, 73(2): 165-174.

Information on priced publications can be obtained from Distribution and Sales (prices in brackets are applicable in developing countries), while copies of documents or off-prints of articles are available free of charge from the Nutrition unit, World Health Organization, 1211 Geneva 27, Switzerland.

Vitamin A deficiency (VAD) is the single most important cause of childhood blindness in developing countries. It also contributes significantly, even at subclinical levels, to morbidity and mortality from common childhood infections. VAD is the result of two primary factors: persistent inadequate intake of vitamin A that is frequently exacerbated by others dietary circumstances, and a high frequency of infections. An estimated 2.8 million preschool-age children are at risk of blindness from VAD, and the health and survival of 251 million others are seriously compromised.

Heightened awareness of the role of vitamin A in human health has led to an international effort to eliminate vitamin A deficiency and its consequences as a public health problem by the year 2000. This is among the important end-of-decade micronutrient goals endorsed by the World Summit for Children (1990), the International Conference on Nutrition (1992), and the World Health Assembly (1993).

This document provides a baseline which national and international health authorities can use to track progress achieved towards the virtual elimination of VAD as a public health problem. It is divided into four sections. The first describes the nature of VAD and reviews the epidemiological issues involved in measuring and interpreting VAD prevalence studies. The second section presents summary tables of the most recent prevalence data, by country and WHO region. The third section presents more detailed sub-national prevalence data by WHO region. The fourth section provides complete bibliographic information for all data sources presented.